



HI-TECH PHARMACEUTICALS

BENZEDRINE™

DIETARY SUPPLEMENT 555 MG

• STRONGEST DIET & ENERGY AID ON THE MARKET!
• THE FIRST ANORECTIC SUPPLEMENT EVER PRODUCED!
• ENHANCES NOREPINEPHRINE, DOPAMINE & SEROTONIN!

60CT

Supplement Facts

Serving Size: 1 Tablet Servings Per Container: 60

Amount Per Serving	Daily Value %
Proprietary Blend with Acacia Rigida 90% Extract (leaves) Yielding the Following Alkaloids as Theano-Ruc: 235mg	
5-Methoxytryptamine HCl, L-Tyrosine Ethyl Ester HCl, Phenylethylamine, N-Benzyl-Alpha-Phenylethylamine	
Proprietary Blend: 100mg	
Theobromine, Alpha-Phenylethylamine HCl, Synephrine HCl, 5-Methoxytryptamine HCl, L-Tyrosine Ethyl Ester HCl, Yohimbe Extract (bark)	
Caffeine (Anhydrous): 200mg	

* Daily value not established

Other Ingredients: Dextrose, Microcrystalline Cellulose, Stearic Acid, Magnesium Stearate, Sodium Stearoyl Glucoside, Starch, Titanium Dioxide, FD&C Blue #2, FD&C Yellow #6, FD&C Red #40, Silica.
Directions: Take 4-2 tablets in the morning and 1 tablet after lunch. Do not exceed 4 tablets daily. Bensedrine is a heat and moisture sensitive, and the bottle should remain sealed after using. Keep dry, and the bottle should remain sealed after using. Temperature 50° - 85°F protected from moisture, heat, and light.

Manufactured by: Hi-Tech Pharmaceuticals, Inc.

6015-B Unity Drive • Norcross, GA 30071 • 1-888-855-7919

THIS PRODUCT HAS NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION.

THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE OR PREVENT ANY DISEASE.

WARNING: NOT FOR USE BY INDIVIDUALS UNDER THE AGE OF 18 YEARS. DO NOT USE IF YOU ARE PREGNANT OR NURSING. INDIVIDUALS WHO CONSUME CAFFEINE WITH THIS PRODUCT MAY EXPERIENCE SERIOUS ADVERSE HEALTH EFFECTS. INDIVIDUALS WHO ARE SENSITIVE TO THE EFFECTS OF CAFFEINE SHOULD CONSULT A LICENSED HEALTH CARE PROFESSIONAL BEFORE CONSUMING THIS PRODUCT.

(WARNINGS) INFORMATION CONTAINED ON THE BACK OF THE LABEL.) SALE TO PERSONS 17 YEARS OF AGE OR YOUNGER IS PROHIBITED.



8 57084 00021 7

Tamper Evident!
Do Not Use if Printed Seal is
Cut, Torn or Missing

IN CASE OF ACCIDENTAL OVERDOSE, SEEK PROFESSIONAL ASSISTANCE OR CONTACT POISON CONTROL CENTER IMMEDIATELY. AVOID ALCOHOL WHILE TAKING THIS PRODUCT. DO NOT EXCEED RECOMMENDED SERVING. EXCEEDING RECOMMENDED SERVING MAY CAUSE SERIOUS ADVERSE HEALTH EFFECTS, INCLUDING HEART ATTACK AND STROKE. DISCONTINUE USE AND CALL A PHYSICIAN OR LICENSED QUALIFIED HEALTH CARE PROFESSIONAL IMMEDIATELY IF YOU EXPERIENCE RAPID HEARTBEAT, DIZZINESS, SEVERE HEADACHE, SHORTNESS OF BREATH, OR OTHER SIMILAR SYMPTOMS. IMPROPER USE OF THIS PRODUCT MAY BE HAZARDOUS TO A PERSON'S HEALTH. EXCEEDING RECOMMENDED SERVING WILL NOT IMPROVE RESULTS. DO NOT USE FOR MORE THAN 8 WEEKS. CONSULT WITH YOUR PHYSICIAN PRIOR TO USE IF YOU HAVE A MEDICAL CONDITION, INCLUDING BUT NOT LIMITED TO: HEART, LIVER, KIDNEY OR THYROID DISEASE, PSYCHIATRIC OR EPILEPTIC DISORDERS, DIFFICULTY URINATING, DIABETES, HIGH BLOOD PRESSURE, CARDIAC ARRHYTHMIA, RECURRENT HEADACHES, ENLARGED PROSTATE OR GLAUCOMA. CONSULT WITH YOUR PHYSICIAN PRIOR TO USE IF YOU ARE TAKING MEDICATION, INCLUDING BUT NOT LIMITED TO: MAOI INHIBITORS, ANTIDEPRESSANTS, ASPIRIN, NONSTEROIDAL ANTI-INFLAMMATORY DRUGS OR PRODUCTS CONTAINING PHENYLEPHRINE, EPHEDRINE, PSEUDOEPHEDRINE, OR OTHER STIMULANTS. DISCONTINUE USE TWO WEEKS PRIOR TO SURGERY.

To report adverse effects call FDA's MedWatch at 1.800.332.1088