## Directions for use:

1 tablet once or twice daily to help promote intestinal health<sup>†</sup>. Dissolve in mouth or chew.

CAUTION: Diff-Stat® should be taken I hour before or 2 hours after antifungal medication.

Do not use if you have a yeast allergy

Keep bottle tightly closed. Store in a cool, dry place.


Do not expose to excessive heat. No refrigeration required. Room temperature stable.

Manufactured in an NSF® GMP registered facility.

Patent Applied For.

<sup>†</sup>These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN.



## Serving Size: 1 chewable tablet

SUPPLEMENT FACTS

FOS, (Fructo-oligosaccharides),

Saccharomyces boulardii, Bacillus coagulans (includes 8.5 billion CFU)

\*Percent Daily Values are based on a 2,000-calorie diet.

\*\*Daily Value not established.

Other ingredients: Sugar, Dextrose, Hydrogenated Cottonseed Oil, Natural & Artificial Orange flavor, Microcrystalline Cellulose, Natural & Artificial Prineapple flavor, Magnesium Stearate, Silicon Dioxide, Ethyl Maltol.

Formulated and marketed by:

Medical Nutrition USA, Inc.

Englewood, NJ 07631 USA
1-800-221-0308 www.diff-stat.info
Copyright ©2010 Medical Nutrition USA. Inc.