

60258-190-01

PRENATABS FA

Dietary Supplement

For Professional Use Only

CYPRESS

PHARMACEUTICAL, INC.

100 Tablets

DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Supplement Facts

Serving Size: 1 Tablet

Amount per serving		%DV*
Vitamin A (Beta Carotene)	4000 IU	50%
Vitamin C (Ascorbic Acid)	120 mg	200%
Vitamin D-3 (Cholecalciferol)	400 IU	100%
Vitamin E (dl-Alpha Tocopheryl Acetate)	30 IU	100%
Vitamin B-1 (Thiamine Mononitrate)	3 mg	176%
Vitamin B-2 (Riboflavin)	3 mg	150%
Niacinamide (Vitamin B-3)	20 mg	100%
Vitamin B-6 (Pyridoxine HCl)	3 mg	120%
Folic Acid	1 mg	125%
Vitamin B-12 (Cyanocobalamin)	8 mcg	100%
Calcium (Calcium Carbonate)	200 mg	15%
Iron (Ferrous Fumarate)	29 mg	161%
Iodine (Potassium Iodide)	150 mcg	100%
Zinc (Zinc Oxide)	15 mg	100%

* Daily Value for Pregnant and Lactating Women.

Other Ingredients: Dicalcium Phosphate, Croscarmellose Sodium, Stearic Acid, Maltodextrin, Crospovidone, Hypromellose, Starch, Sucrose, Citric Acid, Acacia, Titanium Dioxide, Microcrystalline Cellulose, Sodium Lauryl Sulfate, Triacetin, Magnesium Stearate, Magnesium Silicate, Mineral Oil, Polyvinylpyrrolidone, Sodium Ascorbate, Silica, Tocopherols, Tricalcium Phosphate, Triglycerides, FD&C Red #40 Lake and FD&C Yellow #6 Lake.

DIRECTIONS: One tablet daily or as directed by a physician.

CONTRAINDICATIONS: Prenatabs FA is contraindicated in patients with a known hypersensitivity to any of the components of this product. Hemochromatosis and hemosiderosis are contraindications to iron therapy.

WARNINGS:

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of the reach of children. In case of accidental overdose, call a doctor or Poison Control Center immediately.

Folic acid alone is improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where Vitamin B-12 is deficient.

PRECAUTIONS: Folic acid in doses above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations remain progressive. Pernicious anemia should be excluded before using this product since folic acid may mask the symptoms of pernicious anemia.

ADVERSE REACTIONS: Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

STORAGE: Store at room temperature, USP.

Keep out of the reach of children.

To report a serious adverse event or to obtain product information, contact (800) 856-4393.

Distributed By: Cypress Pharmaceutical, Inc., 135 Industrial Blvd., Madison, MS 39110

**NO
VARNISH
HERE**

L353
Rev. 01/11

