

PRE-WORKOUT W/MPE

## **EXPLOSIVE** WORKOUTS

CONTAINS CREATINE NITRATE

HIGHLY EXPLOSIVE ENERGY\* NO3 PUMP TECHNOLOGY\* **RAPID ABSORPTION SYSTEM\*** 







DIETARY SUPPLEMENT | Net. Wt. 342a



	Amount Per Serving	% DV
Calories	5	
Total Carbohydrates	1g	<1%**
Vitamin C	250mg	417%
Calcium	25mg	3%
Niacin (as Niacinamide)	) 30mg	150%
Folic Acid	250mcg	62%
Vitamin B12	35mcg	588%
Beta Alanine	1500mg	t
Creatine Nitrate	1000mg	t
Arginine AKG	1000mg	1
Explosive Energy Blend	741mg	Ť
	N-Acetyl-L-Tyrosine, Caffeine Ani tandardized for L-Dopa). Bitter or	

aurantium) (fruit) (30% Synephrine) (Advantra Z®), Niacinamide, Folate (as Folic Acid), Pyridoxal 5-Phosphate, Vitamin B12 (as Methylcobalamin) Percent Daily Values (% DV) are based on a 2,000 calorie die

Other Ingredients: Natural and Artificial Flavors, Citric Acid, Malic Acid, Silicon Dioxide Calcium Slicate, Sucraiose, Artificial Color (FIBAC Yellow #5), Acesulfame Potassium (Ace-K) Advantra Z@ is a registered trademark of Nutratech Inc. under U.S. patents 6,224,873, 6,316,499, 6,340,481, 6,340,482

SUGGESTED USE:

DO NOT EXCEED RECOMMENDED DAILY INTAKE. USE ONLY AS OIRECTED. Directed Use on Training Days: To determine blerance, begin by taking one serving (1 scoop) mixed with 45 scoop) 20-30 mixed with 45 scoop) 20-30 mixed with 45 scoop) 20-30 mixed serving the serving to the scoop) 20-30 mixtures before training, Attra personal blerance has been assessed, take one to two servings (1-2 scoop) 20-30 mixtures before training begins. Add (4-6 c2) of water for east serving, During your workoul, filts recommended that you drink plenty date. Most serving the serving that serving the serving the serving that serving the serving the serving that serving the serving that serving the servin

\*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.



or any stimulants from other sources whatsoever including but not timited to, coffee, tea, soda and other clietary supplements or medications containing phenoprine or cafeine. Discontinue use and consult your health care professional 2 weeks prior to surgery, or if you experience rapid heartbeat, dizziness, severe headache,



# Supplement Facts

Serving Size: 1 scoop (5.7g) Servings Per Container: 60

	<b>Amount Per Serving</b>	% DV
Calories	5	
Total Carbohydrates	1g	<1%**
Vitamin C	250mg	417%
Calcium	25mg	3%
Niacin (as Niacinamide)	30mg	150%
Folic Acid	250mcg	62%
Vitamin B12	35mcg	588%
Beta Alanine	1500mg	†
Creatine Nitrate	1000mg	†
Arginine AKG	1000mg	T
Explosive Energy Blend	741mg	Ť
Vitamin C (as Ascorbic Acid), N		hydrous

(135mg). Mucuna pruriens (Standardized for L-Dopa), Bitter orange (Citrus aurantium)(fruit)(30% Synephrine)(Advantra Z®), Niacinamide, Folate (as Folic Acid), Pyridoxal 5-Phosphate, Vitamin B12 (as Methylcobalamin)

\*\*Percent Daily Values (% DV) are based on a 2,000 calorie diet. † Daily Value not established.

Other Ingredients: Natural and Artificial Flavors, Citric Acid, Malic Acid, Silicon Dioxide, Calcium Silicate, Sucralose, Artificial Color (FD&C Yellow #5), Acesulfame Potassium (Ace-K) Advantra Z® is a registered trademark of Nutratech Inc. under U.S. patents 6,224,873; 6,316,499; 6,340,481; 6,340,482

## SUGGESTED USE:

DO NOT EXCEED RECOMMENDED DAILY INTAKE. USE ONLY AS DIRECTED. **Directed Use on Training Days**: To determine tolerance, begin by taking one serving (1 scoop) mixed with (4-6 oz.) of water 20-30 minutes before training. After personal tolerance has been assessed, take one to two servings (1-2 scoops) 20-30 minutes before training begins. Add (4-6 oz.) of water for each serving. During your workout, it is recommended that you drink plenty of water. Moisture and humidity can cause clumping and discoloration. STORE IN A COOL DRY PLACE.

IMPORTANT DISCLOSURE: This product may contain legal dietary ingredients that are prohibited by Sports and/or cause a false positive for banned or prohibited substances ("PS"). Due to the unique restrictions of amateur and professional sports organizations, the Armed Forces, law enforcement and other agencies and employers, as well as the possibility of legal dietary ingredients causing a false positive for PS, do not take this product if you are subject to PS testing without the approval and/or authorization of the organization, entity, agency or governing body that requires the testing.

\*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.



715 N. Main Street Bryan, TX 77803 USA www.cellucor.com Product Questions Call: 1.866.927.9686

### WARNING:

Do not take if you are currently taking nitrates for chest pain or if you are taking medication used to freat erectile dysfunction such as PDE-5 inhibitors. This product is only intended to be consumed by healthy adults 18 years of age or older. Before using this product consult with your physician if you are pregnant or nursing, or if you are using any prescription or over the counter medication or if you have any pre-existing medical condition including but not limited to: high or low blood pressure, cardiac arrhythmia, stroke, heart, liver, kidney or thyroid disease, psychiatric or epileptic disorders, diabetes, recurrent headaches, enlarged prostate, difficulty urinating, glaucoma, or if you are taking a MAOI (Monoamine Oxidase Inhibitor), antidepressants, aspirin, nonsteroidal anti-inflammatory drugs or products containing phenylephrine, ephedrine, pseudoephedrine, or any other medication. Contains caffeine. Do not use in combination with synephrine, caffeine or any stimulants from other sources whatsoever including but not limited to, coffee, tea, soda and other dietary supplements or medications containing phenylephrine or caffeine. Discontinue use and consult your health care professional 2 weeks prior to surgery, or if you experience rapid heartbeat, dizziness, severe headache, shortness of breath or any other adverse reaction to this product. Do not combine with alcohol. Do not exceed recommended serving. Do not use if safety seal is broken or missing. Do not use for more than 8 weeks. KEEP OUT OF REACH OF CHILDREN.

