BLACKSTONE LABS

DIRECTIONS OF USADE

As a dietary supplement, take one (1) to two (2) capsules per day with food. Use in cycles of 4 weeks taking at least 4 weeks off in between cycles. Post Cycle Therapy (PCT) is recommended after completion of a cycle. Visit our website for additional PCT information and guidance.


WARNING:

WARNING: KEEP OUT OF REACH OF CHILDREN. Do not use this product if you are at risk for or are being treated for high blood pressure, heart, kidney, thyroid or psychiatric disease, anxiety, depresion, seizure disorders, strokes or any other medical conditions. Do not take this product if you are using anti-depressant, MAO/MAOI inhibitors or any other prescriptions drugs. Not for use by women or by any person under the age of 21. Keep away from children. This product may elevate serum hormone levels wich could produce positive results if you are subject to NCAA testing. For use by healthy idividuals only.

DO NOT USE IF SEAL IS DAMANGED OR MISSING SORE IN A COOL , DRY PLACE.

DISCLAIMER

†These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treate, cure, or prevent any disease.



BLACKSTONE LABS

SUPPLEMENT FACTS

Serving Size:1 Capsule Servings Per Container: 60

Amount Per Serving DV%

4-chloro-17a-methyl-androst-1, 4-dine-3, 17b-dion 50 mg

*(DV) Daily Value Based on a 2,000 calorie diet. ** Daily Value Not Established

Other Ingredients: Magnesium Stearate, Silicon Dioxide,

Titanium Dioxide.

Boca Raton, FL 33433 www.BlackStoneLabs.com



MANUFACTURED EXCLUSIVELY FOR:

BLACKSTONE