201910-480 V2

Cautions and Warnings: If you have gastrointestinal lesions /ulcers, bleeding disorders, or are taking anticoagulant / blood thinners, anti-inflammatory or antibiotic medications, or are about to have surgery, consult a healthcare practitioner prior to use. Do not use if you are pregnant or breast-feeding. If you experience nausea, vomiting or hypersensitivity / allergic reaction while taking Wobenzym, discontinue use and consult your healthcare practitioner.

Recommended dose:

Adults: Take 2 tablets twice daily at least 45 minutes before meals with water.

Store in a cool dry place.
Do not use if safety seal is broken or missing.
Keep out of reach of children.

Distributed by: Mucos USA 112 Technology Drive Pittsburgh, PA 15275 Wobenzym.com 1-800-245-4440



Dietary Supplement



Helps to maintain healthy joints, mobility and flexibility¹

Temporary relief from everyday aches, pains and muscle soreness¹

Speedy recovery following exertion¹



1 These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

Supplement Facts

Serving Size 2 Tablets • Servings per container 240

	Amount Per Serving	% Daily Value
Bromelain** (from pineapple (<i>Ananas comosus</i>) stem)	270 mg (1350 FIP***-ur	* nits)
Trypsin (from porcine (Sus scrofa) or bovine (Bos taurus) pancreas)	144 mg (4320 FIP***-ur	its)
Rutoside trihydrate (Rutin) (from <i>Styphnolobium japonicum</i> flower hud)	300 mg	*

^{*} Daily Value not established.

Other ingredients: Microcrystalline cellulose, calcium phosphate, hydroxypropyl cellulose, vegetable stearate, enteric coating (sodium alginate, medium chain triglycerides, ethylcellulose, stearic acid, oleic acid, purified water) purified water.



^{**} From dried purified aqueous extract.

^{***} FIP is the measurement of enyzme activity according to the methods of the International Pharmaceutical Federation.