Wobenzym[®] N

A Dietary Supplement

Suggested Usage: Take three (3) tablets, two (2) times daily, at least 45 minutes before meals or as recommended by your healthcare professional.

††This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

Warning: If you are pregnant or breast feeding, suffer from bleeding disorders or liver damage, or if you are taking anticoagulants drugs (such as warfarin), consult a healthcare professional before using this product. Store in a cool dry place.

This product does not contain yeast, gluten, soy, milk/dairy, sodium, sugar, artificial coloring or preservatives.

KEEP OUT OF REACH OF CHILDREN DO NOT USE IF SEAL IS MISSING OF RECKEN

200 Tablets Enteric-Coated Manufactured by: Mucos Emulsions GmbH Made in Berlin, Germany Distributed by: 600 Boyce Boad Pittsburgh, PA 15205 USA



Supplement Facts

	Amount Per Serving	% Dail Valu
Pancreatin** 56,000 USP units protease (pancreas) Sus scrofa	300 mg	Ť
Papain** 492 FIP-units*** Carice papaya	180 mg	1
Bromelain** 675 FIP-units Ananas comosus	135 mg	- +
Trypsin** 2,160 FIP-units (pancreas) Sus scrofa and/or Bos taurus	72 mg	†
Chymotrypsin** 900 FIP-units (pancreas) Bos taurus	3 mg	+
Butosida tribudrata** (Butin) Sonbora (anomina	150 mo	+

†Daily Value not established

Other ingredients: cellulose, vegetable-based enteric coating, vegetable

*** FIP is the measurement of enzyme activity according to the methods



Wobenzym®N is backed by decades of clinical research