

Each tablet is in a plastic blister with a foil back.

To remove tablet from blister:

- Separate blister along perforationPeel back at arrow
 - Push tablet through from front





100 TABLETS 325 mg each TAMPER EVIDENT: Do not use if blister units are torn, broken, or show signs of tampering.



Supplement Facts

Serving Size: 1 Tablet

Amount % Daily Per Tablet Value

IRON (as Ferrous Sulfate) 65mg 361%

Each tablet contains 325mg (5gr) Ferrous Sulfate, providing approximately 65mg of elemental iron.

OTHER INGREDIENTS: Cellulose, Crospovidone, Stearic Acid, Sodium Lauryl Sulfate, Magnesium Silicate, Hypromellose, Silica, PEG, FD&C Blue #2 Lake, Mineral Oil, Titanium Dioxide, FD&C Yellow #5 Lake (Tartrazine), FD&C Blue #1 Lake, FD &C Red #40 Lake, FD&C Yellow #6 Lake, Calcium Sulfate.

DIRECTIONS: Adults: take one (1) tablet daily with or after a meal as a dietary supplement, or as recommended by your physician. Do not exceed recommended dosage.

Store at room temperature

REV 0810

WARNING: Accidental overdose of iron containing supplements is a leading cause of fatal poisoning in children under 6.

KEP THIS PRODUCT OUT OF REACH OF CHILDREN.
In case of accidental overdose, seek professional assistance

WARNINGS: The treatment of any anemic condition should be under the advice and supervision of a physician. Since oral iron products interfere with the absorption of oral tetracycline

or contact a poison control center immediately.

antibiotics, these products should not be taken within two hours of each other.

As with any supplement, if you are pregnant, pursing, or

As with any supplement, if you are pregnant, nursing, or taking medication, ask a health professional before use.

This product may occasionally cause constipation, diarrhea, or may color stool black. It may cause gastrointestinal discomfort (such as nausea) which may be minimized by taking with meals.

DISTRIBUTED BY: Geri-Care Pharmaceuticals Corp. 1650 63rd Street Brooklyn, NY 11204 3 57896 70301