## SUPPLEMENT FACTS

Serving Size: One (1) Capsule

Servings Per Container: 60

## **Amount Per Serving**

% Daily Value\*\*

19-Norandrosta-drol TM 65mg\*\*

3-beta-Hydroxy-5-androsten-17-one

1, 3, 7-trimethyl-1 H-purine-2,6(3H, 7H)-dione

2-amino-5-(diaminomethylidene-amino) pentanoic acid

17a-di methyl-bol TM 45mg\*\*

3beta-hydroxyandrost-5-ene-17-one 4-aminobutanoic acid

\*\*Daily Value Not Established

OTHER INGREDIENTS: Gelatin, Silica, Magnesium Stearate, Maltodextrose, Cellulose. Warning: CONSULTYOUR PHYSICAIN BEFORE USING THIS PRODUCT. Warning: NOT FOR WOMEN OR INDIVIDUALS UNDER THE AGE OF 18 YEARS. This product is only intended to be consumed by healthy adult males 18 years of age or older. Consult a physician or licensed qualified health care professional before using this product if you have any pre-existing medical condition including but not limited to: high or low blood pressure, cardiac arrhythmia, anxiety, depression, stroke, heart, liver, kidney or thyroid disease, seizure disorder, psychiatric disease, diabetes, prostate cancer, prostate enlargement, low "good" cholesterol (HDL), any other medical condition, or if you are using any other dietary supplements, prescription drug, or over-the-counter drug. Do not exceed recommended serving. Exceeding recommended serving may cause serious adverse health affects. Possible side affects include acne, hair loss, hair growth on the face (in women), aggressiveness, irritability, and increased levels of estrogen. Discontinue use and call a physician or licensed qualified health care professional immediately if you experience rapid heartbeat, dizziness, blurred vision, or other similar symptoms.



If you want crazy gains in mass and strength, then you should check out TOKKYO STACK EXTREME, the strongest double stack supplement available!

**Directions:** As a dietary supplement, take one (1) capsule in the morning and one (1) capsule in the afternoon. For maximum absorption, take this supplement with at least 15 grams of fat. It is recommended that you take Tokkyo Liver-X for the duration of this supplement, and take Tokkyo Clomidex for at least two weeks after discontinuing use of this supplement.

Note: If you are a professional or amateur athlete subject to performance enhancing substance testing, do not use this product unless cleared by your sanctioning body as it may cause a false reactive result for a urine or blood drug test.

\*These statements have not been evaluated by the FDA This product is not intended to diagnose, treat, cure, or prevent any disease.

Allergy Warning: Manufactured in a facility that processes milk, soy and egg proteins, nuts, and other tree nuts and wheat.

READ ALL LABEL INFORMATION AND WARNINGS. DO NOT EXCEED RECOMMENDED DOSAGE.



DISTRIBUTED BY: BTE WHOLESALERS, LLC
587 2ND STREET, SUITE B • MANCHESTER, NH 03102, USA
1.866.320.0141 WWW.TOKKYONUTRITION.COM