For more information, please visit integrative pro.com. THIS STATEMENT HAS NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.

Recommendation: Take 2 tablets at bedtime or as recommended by your healthcare professional. Reduce dosage if you experience abdominal discomfort or diarrhea. If discomfort persists, discontinue use. Do not take more than 2 tablets in a 24-hour period or use the maximum dosage for more than 7 days. Not for long term use.

Keep out of reach of children. Safety sealed with printed inner seal. Do not use if seal is broken or missing.

Contains no sugar, yeast, soy, dairy products, artificial colors, flavors, preservatives, or ingredients of animal origin.

©2018 INTEGRATIVE THERAPEUTICS, LLC GREEN BAY, WI 54311 USA integrativepro.com • 800.931.1709

♠ LZ78416.A01 BLK841A



INTEGRATIVE THERAPEUTICS®

LAXATIVE FORMULA

60 TABLETS

DIETARY SUPPLEMENT

Supplement Facts

Serving Size 2 Tablets Si	ervings per Conta	iner 30
Amount per 2 Tablets		% DV
Total Carbohydrate	1 g	<1%†
Sodium	5 mg	<1%
Magnesium Hydroxide	1.7 g	**
Proprietary Mucilage Blend marshmallow (Althaea officinalis) root extract, slipper elm (Ulmus rubra) bark, and fenugreek (Trigonella foe graecum) seed extract		**
Proprietary Soothing Blend fennel (Foeniculum vulgare) seed, ginger (Zingiber off rhizome extract, and peppermint (Mentha x piperita) li		**

Percent Daily Values (DV) are based on a 2,000 calorie diet. **Daily Value not established

Other ingredients: cellulose, sodium carboxymethyl cellulose, stearic acid, hydroxpropyl methylcellulose, magnesium stearate, glycerin, silicon dioxide

Warning: Do not use this product if pregnant, nursing, attempting to become pregnant, or under the age of 12. Do not exceed recommended dose, or take longer than the recommended period, as frequent or prolonged use may result in dependence. Do not use when abdominal pain, nausea or vomiting are present. Consult your healthcare professional if you notice a sudden change in bowel habits that persist over two weeks. Rectal bleeding or failure to move bowels after use may indicate a serious condition. Discontinue use and consult your healthcare professional. This product contains Slippery Elm. Slippery Elm should not be taken together with any medication as it may interfere with their absorption. If you have diabetes, consult your healthcare professional before use