Unflavored Tablets

Place two tablets in mouth. Let dissolve and swallow. Follow recommended usage table below:

 100-120 Lbs
 2 Tablets 3 times per day

 121-170 Lbs
 2 Tablets 4 times per day

 171-220 Lbs
 2 Tablets 5 times per day

 221 Lbs+
 2 Tablets 6 times per day

Do not exceed recommended use

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

Supplement Facts

Serving Size: 2 Tablets Servings Per Container: 150

Amount Per Serving Value

Vitamin A (as retinyl acetate) 834 IU 16%

Vitamin D3 (as cholecalciferol) 66 IU 16%

Anatabine (as anatabine citrate) Anti-Inflammatory Support*

Dietary Supplement

Anatabloc® is Gluten-Free

† Daily Value not established.

Other Ingredients: Mannitol, silicon dioxide, stearic acid, hydroxypropyl methylcellulose, magnesium stearate, titanium dioxide (color), natural

2 mg †

Other Ingredients: Mannitol, silicon dioxide, stearic acid, hydroxypromethylcellulose, magnesium stearate, titanium dioxide (color), natural glaze, methylparaben, propylparaben, triethyl citrate, citric acid, BHT, mono and diglycerides, and polysorbate 80.

For optimum freshness keep the product cool and dry.



Manufactured for Rock Creek Pharmaceuticals 55 Blackburn Center, Gloucester, MA 01930 Learn more at: anatabloc.com



Unflavored Tablets

Anatabloc

Anti-Inflammatory Support*
Dietary Supplement

50%
MORE

Anatabloc

Unflavored Tablets

Anti-Inflammatory Support*
Dietary Supplement



Anatabloc

Unflavored Tablets

Anti-Inflammatory Support*
Dietary Supplement

50% MORE