

					The second second
upplen ing Size ings Per Container	1/2 Tablet 160	Facts 1 Tablet 80		% DV for Children 2 and 3 years (1/2 Tablet)	% DV for Children 4 years and older (1 Tablet)
	% DV for Children 2	% DV for Children	Folic Acid 400 mcg	100%	100%
	and 3 years	4 years and	Vitamin B ₁₂ 6 mcg	100%	100%
ount Per Tablet	(1/2 Tablet)	older (1 Tablet)	Biotin 45 mcg	15%	15%
ories 5			Pantothenic Acid 10 mg	100%	100%
Carbohydrate 1 g	*	<1%†	Calcium 108 mg	7%	11%
gars <1 q	*	*	Iron 18 mg	90%	100%
min A 3.500 III	70%	70%	Phosphorus 50 mg	3%	5%
% as Beta-Carotene	e) (9	1070	lodine 150 mcg	107%	100%
min C 60 mg	75%	100%	Magnesium 40 mg	10%	10%
min D 400 IU	50%	100%	Zinc 15 mg	94%	100%
min E 30 IU	150%	100%	Copper 2 mg	100%	100%
min K 10 mcg	*		Manganese 1 mg	*	50%
min 1.5 mg	107%	10/0	Chromium 20 mcg	*	17%
flavin 1.7 mg	106%	10070	Molybdenum 20 mcg	*	27%
nin B6 2 mg	111%	1000/		tablished	
	143%	100%	* Daily Value (DV) not es † Percent Daily Values ar	o based on a 2	2.000 calorie diet.
lients: Sucrose Da		10070	Percent Daily Values at	C Dascu on a	Coloium Carbonate

Acid (Vit. C), Magnesium Oxide, Pregelatinized Corn Starch, Mono- and Di-glycerides, Stearic Acid Maltodextrin. Contains < 2% of: Aspartame,** Beta-Carotene, BHT (to retard oxidation), Biotin, Blue 2 dicum Pantothenate, Carbonyl Iron, Carrageenan, Cholecalciferol (Vit. D3), Chromic Chloride, Citric Acid, Cyanocobalamin (Vit. B12), Dextrose, dI-Alpha Tocopheryl Acetate (Vit. E), Folic Acid, Gelatin, Guar Gravors, Niacinamide, Phytonadione (Vit. K), Potassium Iodide, Pyridoxine Hydrochloride (Vit. B6), Red 40 doction (Vit. B2), Silicon Dioxide, Sodium Molybdate, Thiamine Mononitrate (Vit. B1), Tocopherols (to retard oxidation), Vitamin A Acetate, Vellow Col.

Use: Children 2 and 3 years of age, chew approximately 1/2 tablet daily with food. Children 4 years of age and old tablet daily with food. Not formulated for use in children less than 2 years of age. Do not exceed suggested use.

RNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6, and this product out of reach of children. In case of accidental overdose, call a doctor or poison control center increase.

"PHENYLKETONURICS: CONTAINS PHENYLALANINE.

MPORTANT INFORMATION: Long-term intake of high levels of vitamin A (excluding that sourced from beta-carotene) may increase the risk of osteoporosis in adults. Do not take this product if taking other vitamin A supplements.

Some at room temperature. Keep bottle tightly closed. Protect from moisture.

and sealed with printed foil under cap. Do Not Use if foil is torn.

Marketed by: Pfizer, Madison, NJ 07940 USA © 2011 Pfizer Inc. Made in Canada

Questions? Comments? Call 1-877-CENTRUM

most recent product information, visit www.centrum.com

neters to key nutrients for growth and development.



PAA024517

CONTROL NO.

F63615

EXP. DATE

tally with food. Children 4 years of age and older,

Marke in Canada

e of fatal poisoning in children under 6.
a doctor or poison control center immediately.

A jexcluding that sourced from beta-carotene) may increase

PAA024517

CONTROL NO.

F63615 NOV 13

EXP. DATE