TIRECTIONS As a dietary supplement,

take one (1) to two (2) capsules per day with food . Use in cycles of 4 weeks taking at least 4 weeks off in between cycles. Post Cycle Therapy (PCT) is highly recommended after completion of a cycle. Visit our website for additional PCT information and guidance.

DO NOT USE IF SEAL IS DAMAGED OR MISSING. STORE IN A COOL, DRY PLACE

Do not exceed the recommended dose. Keep out of reach of children. Consult your physician before using this dietary supplement. Do not use this product if you are at risk or are being treated for high blood pressure, heart, kidney, thyroid, or psychiatric disease, difficulty in urinating, prostate enlargement, seizure discorder, or stroke.

†These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose. treat, cure, or prevent any disease.

BLACKSTONE -----LABS-



SUPPLEMENTS FACTS

Serving Size: 1 Capsule Servings per Container: 60

Amount per serving

Estra-4.9.11-triene-3.17-dione

% DV * 15ma*

* (DV) Daily Value Based on a 2,000 calorie diet * * Daily Value not established

Other Ingredients: Magnesium Stearate, Silicon Dioxide, Titanium Dioxide



BLACKSTONE CHOYALABS

Boca Raton, FL 33433

www.BlackStonel.abs.co.