STORAGE: Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F). Brief exposure to temperatures up to 40°C (104°F) may be tolerated provided the mean kinetic temperature does not exceed 25°C (77°F); however, such exposure should be minimized.

NOTICE: Contact with moisture can discolor or erode the soft gel.

HOW SUPPLIED: Bottles of 30 soft gels

each - NDC 0178-0896-30.

To report a serious adverse event or obtain product information, call (800)

298-1087.

Please consult your health care

Please consult your health care provider with any dietary concerns.

citranatal.com

Trademark of Mission Pharmacal Company



CitraNatal Medley contains We'sDHA®, a trademark of DSM

CitraNatal Medley soft gels are MADE IN CANADA for: MISSION PHARMACAL COMPANY San Antonio, TX IISA 78230 1355

Mission®

Copyright © 2018 Mission Pharmacal Company.

All rights reserved. L089630R0618



WARNING: Accidental overdose of

cause of fatal poisoning in children

poison control center immediately.

iron-containing products is a leading

under 6 KEEP THIS PRODUCT OUT

OF THE REACH OF CHILDREN, In case

of accidental overdose, call a doctor or

NDC 0178-0896-30

iron-containing products is a leading cause of fatal poisoning in children under 6. KEP THIS PRODUCT OUT OF THE REACH OF CHILDREN. In case of accidental overdose, call a doctor or poison control center immediately.

WARNING: Ingestion of more than

Rx Only WARNING: Accidental overdose of

3 grams of omega-3 fathy acids per day has been shown to have potential antithrombotic effects, including an increased bleeding time and INR. Administration of omega-3 fathy acids should be avoided in patients on anticoapylarits and in those known to have an inherited or acquired bleeding diathesis. WARNING: Folic acid alone is improper therapy in the treatment of permicious amenia and other megaloblastic

anemia and other megaloblastic anemias where vitamin B₁₂ is deficient. L089630R0618

DESCRIPTION: CitraNatal MedigyTM is a prescription prenatalpostantal multivitamin'mineral soft get with Ferr-Ease[®], a patented dual-ien delivery comprising both a quick release and slow release iron, and DHA an essential fattly acid. The prenatal vitamin is an oval, purple, opaque soft get containing a greenish-gray liquid to semi-solid fill. The prenatal soft cel is printed "599" in white ink.

Each prenatal soft gel contains: Vitamin D₃ (cholecalciferol) 200 IU

 Vitamin E (dl-alpha tocopheryl acetate)
 15 IU

 (dl-alpha tocopheryl acetate)
 12.5 mg

 Folic Acid
 1 mg

 Calcium (calcium citrate)
 62 mg

 Iron (carbonyl iron, ferrous fumarate)
 27 ma

Docosahexaenoic Acid (DHA, 53% from 377 mg algal oil)...... 200 mg

Inactive Ingredients: Gelatin, glycerin, lecithin, titanium dioxide, ethyl vanillin, D&C Red #33, butylated hydroxytoluene (BHT), FD&C Blue #1, and white ink.

INDICATIONS: CitralNatal Medley is a multi-vitamin/mineral prescription drug indicated for use in improving the nutritional status of women prior to conception, throughout pregnancy, and in the postnatal period for both lactating and nonlactating mothers.

CONTRAINDICATIONS: This product is contraindicated in patients with a known hypersensitivity to any of the ingredients.

WARNING: Ingestion of more than 3 grams of omega-3 fatty acids per day has been shown to have potential antithrombotic effects, including an increased bleeding time and INR. Administration of omega-3 fatty acids should be avoided in patients on anticoacquiants and in those known

bleeding diathesis.

WARNING: Folic acid alone is improper therapy in the treatment of pernicious

to have an inherited or acquired

anemia and other megaloblastic anemias where vitamin B₁₂ is deficient. **PRECAUTIONS:** Folic acid in doses above 0.1 mg daily may obscure

above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations progress.

ADVERSE REACTIONS: Allergic

ADVERSE REACTIONS: Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

CAUTION: Exercise caution to ensure that the prescribed dosage of DHA does not exceed 1 gram (1000 mg) per day.

DOSAGE AND ADMINISTRATION:

One prenatal soft gel daily or as directed by a physician.