

Suggested Use: As a dietary supplement, take 4 tablets per day with food. To accommodate body weight, activity level, stress level, and/or inadequate diet, take additional tablets up to a maximum of 8 tablets per day.

Contains no artificial colorings, fillers, sweeteners, fragrance or preservatives. Also contains no yeast, milk, egg, gluten, starch or wheat. This product is not a substitute for any medication. If you are pregnant, nursing, or suffer from a medical condition, consult a physician before taking this product.

Another quality supplement from **FACTOR**
NUTRITION LABS

*A recent clinical study showed that adults who took FOCUSfactor as directed for six weeks improved their memory, concentration, and focus.

**This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

#476480



LOT# 50029265
EXP. 03/2017

AMERICA'S #1 BRAIN HEALTH SUPPLEMENT

FOCUSfactor

Nutrition for the Brain

Clinically Shown* to Improve
MEMORY, CONCENTRATION and FOCUS**



DIETARY SUPPLEMENT

150 Tablets

Supplement Facts

Serving Size: 4 tablets

	Amount Per Serving	% DV
Vitamin A (from natural beta-carotene and mixed carotenoids)	4,000 IU	80%
Vitamin C (as ascorbic acid, sodium ascorbate, calcium ascorbate, potassium ascorbate, zinc ascorbate, and ascorbyl palmitate)	250 mg	417%
Vitamin D (as cholecalciferol)	100 IU	25%
Vitamin E (as natural d-alpha-tocopheryl succinate)	30 IU	100%
Thiamin (as thiamin mononitrate)	3 mg	200%
Riboflavin	1.7 mg	100%
Niacin (from inositol hexanicotinate and 50% as niacinamide)	25 mg	125%
Vitamin B6 (as pyridoxal-alpha-ketoglutarate, pyridoxal 5-phosphate)	15 mg	750%
Folate (as folic acid)	400 mcg	100%
Vitamin B12 (as ion exchange resin)	20 mcg	333%
Biotin	300 mcg	100%

Synergistic and proprietary formulation

Dimethylaminoethanol (as DMAE bitartrate), L-glutamine, Bacopin® (bacopa monnieri extract; leaf)†††, L-pyrroglutamic acid, phosphatidylserine, docosahexaenoic acid concentrate (15% DHA from fish (anchovy, sardine, and salmon) body oil, phosphatidylcholine, inositol, N-acetyltyrosine, bitter melon fruit standardized extract (25% anthocyanosides), gamma-choline (as choline bitartrate), inositol, N-acetyltyrosine, bitter melon fruit standardized extract (25% anthocyanosides), gamma-aminobutyric acid, grape skin extract and Activin™ (grape seed extract)††††, vinpocetine, Trace-Lyte™ electrolyte concentrate, huperzine A (extract of huperzia serrata; whole plant), boron (as boron citrate) and vanadium (as vanadyl sulfate).

*Daily Value not established.

Other ingredients: Dicalcium phosphate, microcrystalline cellulose, croscarmellose sodium, stearic acid, silica, magnesium stearate and pharmaceutical glazes. †Bacopin® is a trademark of the ISP Group. ††Bacopin® is a registered trademark of Albion Laboratories, Inc. †††Bacopin® is a registered trademark of Sabinsa Corporation. ††††Activin™ is a trademark of San Joaquin Valley Concentrates.

Store at 15-25°C (59-77°F). • Protect from heat, light and moisture. • Do not purchase if seal is broken.

Distributed by Factor Nutrition Labs, LLC
865 Spring St. Westbrook, Maine 04092
www.factornutrition.com • 1-800-515-2143

U.S. Patent Pending • Made in U.S.A.
Tracking Code: FOCUSF010150K51 / 2205878

Servings Per Container: 37.5

	Amount Per Serving	% DV
Pantothenic acid (as d-calcium pantothenate)	12 mg	120%
Calcium (as calcium citrate, calcium succinate and calcium ascorbate)	50 mg	5%
Iron (as Ferronyl™ carbonyl iron)††	5 mg	28%
Iodine (as kelp (ascophyllum nodosum))	15 mcg	10%
Magnesium (as magnesium citrate, magnesium malate and magnesium taurinate)	100 mg	25%
Zinc (as zinc citrate and zinc ascorbate)	10 mg	67%
Selenium (as selenomethionine)	50 mcg	71%
Copper (as copper citrate, TRAACS® copper glycinate chelate)†††	0.4 mg	20%
Manganese (as manganese citrate)	2 mg	100%
Chromium (as chromium polynicotinate)	100 mcg	13%
Molybdenum (as molybdenum amino acid chelate)	10 mcg	83%
Potassium (as potassium citrate, potassium aspartate and potassium ascorbate)	50 mg	1%
	692 mg	

*Reference: Journal of the American Medical Association, 2000; 283:2560-2563. **Reference: Journal of the American Medical Association, 2000; 283:2560-2563. ***Reference: Journal of the American Medical Association, 2000; 283:2560-2563. ****Reference: Journal of the American Medical Association, 2000; 283:2560-2563.