



Moloco + B12

Manufacturer	Darya-Varia
Contents	Placental extract, vitamin B12.
Indications/Uses	Primary and secondary hypogalactia.
Dosage/Directions for Use	Three (3) dragees daily (1 dragee after each main meal) during the whole period of lactation ie, during the first 9 months after birth. Moloco + B12 dragees may either be swallowed whole or chewed. Experience has shown that milk flow may stop if cure is interrupted.
Special Precautions	Although many excellent baby foods are available, none of them, not even specially prepared cow's milk, is altogether suitable for infants. Exceptionally strong babies may tolerate them, but the more delicate and weak are subject to constant digestive disturbances. It must be emphasized that the only correct food for a child during the first months of life is its mother's milk. Human milk cannot be replaced or improved by any other kind of food. It is the only natural nutrient for babies. Most of the infants dying within the first 9 months of life are bottle fed.
Description	Each dragee contains placental extract 15 mg and vitamin B12 (cyanocobalamin) 20 mcg.
Mechanism of Action	The hormones and other active components of Moloco + B12 stimulate the activity of the milk glands to such an extent that mothers will be able to breastfeed their children and thus, provide them effective protection against outside influences. Numerous medical experiments have proved the correctness of this assertion and have given convincing evidence that regular ingestion of Moloco + B12 does not cause the slightest inconvenience to mother and child.
MIMS Class	Supplements & Adjuvant Therapy
ATC Classification	G02CX - Other gynecologicals ; Used in the management of gynecological conditions.
Regulatory Classification	G
Presentation/Packing	Dragee (sugar-coated) 60's.