Integra - 30 NOC 52747-710-30 30 Capsules / One a day INTEGRA IRON / VITAMIN SUPPLEMENT CAPSULES INDICATIONS: Integra™ is indicated for the treatment of iron deficiency anemia, which may occur due to an increased need for iron, a deficient intake of iron, or an excessive loss of iron. DOSAGE AND ADMINISTRATION: Adults (persons over 12 years of age), one (1) capsule

daily, between meals, or as prescribed by a physician. Do not exceed recommended dosay

Rev. 02/2015

Do not administer to children under the age of

Store at controlled room temperature 15° to 3 (59° to 86° F). Keep in a cool, dry place. Caps

are non USP.



Other Ingredients: Hypromellose, Magnesium Stearate, Titanium Dioxide, FD & C Red #40, FD & C Blue #1.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

Consult package literature for full prescription information. You should contact your healthcare provider for medical advice about adverse events. To report a serious adverse event, contact US Pharmaceutical Corporation. P.O. Box 360465, Decatur. GA 30036. Marketed by US Pharmaceutical Corporation. Manufactured with Vcaps® Plus capsule shells. Store at controlled room temperature 15° to 30°C (59° to 86° F). Keep in a cool, dry place. Manufactured in a FDA registered facility in the USA.

Iron / Vitamin Supplement - 30 Capsules

Consult package literature for full prescription information. You should contact your healthcare provider for medical advice about adverse events. To report a serious adverse event, contact US Pharmaceutical Corporation, P.O. Box 360465, Decatur, GA 30036. Marketed by US Pharmaceutical Corporation. Manufactured with Vcaps® Plus capsule shells. Store at controlled room temperature 15° to 30°C (59° to 86° F). Keep in a cool, dry place. Manufactured in a FDA registered facility in the USA.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of acciden overdose, call a doctor or poison control center immediately.

US Pharmaceutical Corporation, P.O. Box 360465, Decatur, GA 30036 (USA) http://www.uspco.com