

Prenatal Tablets

*A Dietary Supplement For
Pregnant & Lactating Women*

30 Tablets

Supplement Facts

Serving Size: 1 Tablet

Amount Per Serving		% Daily Value
Vitamin A (as Acetate and Beta Carotene)	4000 I.U.	50%
Vitamin C (as Ascorbic Acid)	100 mg.	167%
Vitamin D-3 (Cholecalciferol)	400 I.U.	100%
Vitamin E (as dl-Alpha Tocopheryl Acetate)	11 I.U.	37%
Vitamin B-1 (as Thiamine Mononitrate)	1.5 mg.	88%
Vitamin B-2 (Riboflavin)	1.7 mg.	85%
Vitamin B-3 (as Niacinamide)	18 mg.	90%
Vitamin B-6 (as Pyridoxine HCl)	2.6 mg.	104%
Folic Acid	0.8 mg.	100%
Vitamin B-12 (as Cyanocobalamin)	4 mcg.	50%
Calcium (as Calcium Carbonate and Calcium Sulfate)	263 mg.	26%
Iron (as Ferrous Fumarate)	27 mg.	150%
Zinc (as Zinc Oxide)	25 mg.	167%

PEEL BACK FOR MORE INFORMATION

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under six. **KEEP THIS PRODUCT OUT OF THE REACH OF CHILDREN.** In case of accidental overdose, call a doctor or Poison Control Center immediately.

STORE IN A DRY PLACE AND AVOID EXCESSIVE HEAT. TAMPER RESISTANT: DO NOT USE IF SEAL UNDER CAP IS BROKEN OR MISSING.

Distributed by:
MAJOR[®] PHARMACEUTICALS
31778 Enterprise Drive
Livonia, MI 48150, USA
Questions or Comments?
(800) 616-2471

M-29 Rev 06/13

OTHER INGREDIENTS:

Cellulose, Stearic Acid, Croscarmellose Sodium, Hypromellose, Gelatin, Acacia, Maltodextrin, Titanium Dioxide, Starch, Triacetin, Magnesium Stearate, Magnesium Silicate, Dicalcium Phosphate, Mineral Oil, Sucrose, Silica, Sodium Ascorbate, Triglycerides, FD&C Red #40 Lake, FD&C Yellow #6 Lake, Mixed Tocopherols, Sodium Benzoate, Sorbic Acid, Tricalcium Phosphate, BHT and Ascorbyl Palmitate.

DIRECTIONS: For adults, take one (1) tablet daily, preferably with a meal. As a reminder, discuss the supplements and medications you take with your health care provider.

FREE OF: Yeast, wheat, milk or milk derivatives, soy, artificial flavor, sodium (less than 5 mg per serving).

WARNING: If you are pregnant, nursing, taking any medications or planning any medical procedure, consult your doctor before use. Discontinue use and consult your doctor if any adverse reactions occur.