

Mix one packet per day with at least 12 oz of water or any non-carbonated beverage. Stir or blend well to completely dissolve.

Use before, during and after pregnancy



Plant-based Omega-3, Ferrochel® Gentle DHA and Choline to Iron to help minimize support fetal brain development."

iron-induced constipation."



B Vitamins to Contains Folate, the boost energy, May necessary ingredient in prenatal vitamins also help with for early fetal morning sickness." development."

## drinkpremama.com

\*These statements have not been evaluated by the Food and Drug Administration.

Ferrochel is a registered trademark of Albion Laboratories, Inc.



Prenatal Drink Mix

> with DHA Citrus Flavored Multivitamin Supplement

> > STAGE 3

GLUTEN FREE . VEGETARIAN . NON-GMO NET WT 3.7 OZ (106g)

ner	١t	Fa	ct	s	
acket	13.8	a)			

Amount Per Serving	100	% Daily Velo for Prognat & Leptation Woman		
Calories	5			
Total Cartiohydrate	1.59	1%		
Vitamin C (as ascorbic acid, USP-FCC)	30mg	25%		
Vitamin D (as Vitamin D2)	22.5 mog (900 fU)	150%		
Ribofavin (USP-FCC)	1.70	10674		
Vitamin B6 (as pyridexine hydrochloride, USP-FCC	30mg	1500%		
Paint.	143mog DFE og Folic Acid)	190%		
Vitamin B12 (as cyanonchalamin, USP)	6mog	214%		
Biotin (USP)	300mog	857%		
Choline (as choline bitartrade, USP-FCC)	25mg	5%		
Calcium (as calcium factate, USP-FCC and dicalcium phosphate, FCC)	100mg	8%		
ron (as ferrous pyrophosphate encap)	9mg	33%		
Magnesium (as magnesium citrate, USP)	60mg	157		
Zinc (as zinc citrate, USP)	7.5mg	581		
Copper (as copper gluconate, USP)	2mg	1547		
Omega-3 Fathy Acid Bland	200mg	1		
CHA (docosahexaenois acid) (as algal oil)	20mg	1		

Other Ingredients: Natural Flavors, Citric Acid, Stevia Extract, Talin

CONTAINS TRACES OF MILK AND SOY WARRING: Accidental overdose of iron-containing products is a leading cause of fulfill

poisoning in children under the age of 6. Keep this product out of reach of children in case of accidental overdose, call a doctor or polson control center immediately As with any supplement, if you are pregnant, breastfeeding or taking medication, consult your doctor before use.

Store at room temperature. Avoid excessive heat. KEEP OUT OF REACH OF CHILDREN.

Do not use this product if safety seal on flap is torn of