

Important: Use Axis-HT for at least 6 weeks in order to fully maximize its anabolic-androgenic potential. Do not exceed 12 weeks of continuous use. NOTE: If a reduction in androgenic effects is desired, cut dosage in half.

SUPPLEMENT FACTS

Serving Size: 2 Tablets
Servings Per Container: 60

	Amount Per Serving	% Daily Value
Magnesium (magnesium aspartate)	150mg	38%
Zinc (monomethionine and aspartate)	10mg	67%
StAR-P2™ (Patent Pending)	2238mg*	
Androgenic-Anabolic Testosterone Matrix		
Inositol Hexaphosphate		
TPS3™ (3 Fraction Tribulus Terrestris composite yielding a precision ratio of Protodioscin and Saponin concentrations)		
GF5™ (Proteinaceous Avian Nucleus Extract containing naturally occurring growth factors: IGF, FGF, NGF, EPF, GTCF)		
Testofen™ (Fenugreek Extract Standardized for 50% Fenuside)		
Safed Musli		
Eurycoma Longifolia		
Coleus Forskohlii		
(Standardized for 40% Forskolin)		

* Daily Value Not Established

Other Ingredients: Microcrystalline Cellulose, Stearic Acid, Povidone, Hydroxypropyl Methyl Cellulose, & Polyethylene Glycol



Pro-Testosterone Amplifier*

**A Designer Anabolic
& Androgenic Potentiator***

RECOMMENDED USE: As a dietary supplement, take 2 tablets 3 times per day, everyday. Take the first dose in the morning with breakfast, the second with lunch, and the third at night with your last meal. Axis-HT works synergistically with protein and carbohydrates to help promote muscle anabolism. It is therefore highly recommended that you take Axis-HT with a protein and carbohydrate-rich meal or shake. Stack Axis-HT with NO-Xplode, CellMass, and Nitrix.

WARNING: Seek advice from a health care practitioner prior to use if you have any pre-existing medical condition including, but not limited to: hypertension, hypotension, heart, liver, or thyroid disease, prostate hypertrophy, diabetes, psychological disorders, asthma, pernicious anemia, anxiety, depression, seizure disorder, cardiac arrhythmia, stroke, difficulty urinating due to prostate enlargement, or if you are taking an MAO inhibitor or any other prescription drug. Do not use if you are pregnant or nursing. Reduce or discontinue use if excessive sleeplessness, tremors, dizziness, nervousness, headaches, or heart palpitations occur. For use by healthy adults only. Not intended for use by persons under the age of 18. Keep out of reach of children. After 12 weeks of use discontinue for at least 4 weeks.

Customer Inquiries: Call 1-877-673-3727
Developed By & Manufactured Exclusively For:
Bio-Engineered Supplements & Nutrition
Boca Raton, FL 33487 USA



*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.