

51991-203-11

Ferrex™ 150

Polysaccharide-Iron Complex Capsules

**DO NOT USE IF BLISTER
SEALS ARE BROKEN**

breckenridge | A Towia
Company

Dietary Supplement
100 Unit Dose Capsules

Supplement Facts

Serving Size: 1 Capsule
Servings Per Container: 100

	Amount Per Capsule	% Daily Value
Iron (Elemental) (as Polysaccharide-Iron)	150 mg	833%

Other ingredients: Sodium Lauryl Sulfate, Gelatin, Stearic Acid, Polyethylene Glycol, Microcrystalline Cellulose, Citric Acid, Croscarmellose Sodium, Titanium Dioxide, Polyvinylpyrrolidone, FD&C Yellow #5 Lake*, FD&C Red #40 Lake, FD&C Red #3 Lake and FD&C Blue #1 Lake.

* Contains FD&C Yellow No. 5 Lake (Tartrazine) as a color additive. (see **PRECAUTIONS**)

Ferrex™ 150 Polysaccharide-Iron Complex Capsules are a product of ferric iron complexed to a low molecular weight polysaccharide.

INDICATIONS AND USAGE

Ferrex™ 150 may be used for the dietary management of iron deficiencies.

CONTRAINDICATIONS

Ferrex™ 150 is contraindicated in patients with a known hypersensitivity to any of the components of this product. Hemochromatosis and hemosiderosis are contraindications in the use of this dietary supplement.

PRECAUTIONS

This product contains FD&C Yellow No. 5 Lake (Tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No. 5 Lake (Tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

The type of anemia and the underlying cause or causes should be determined before initiating supplementation with Ferrex™ 150. Since the anemia may be a result of a systematic disturbance, such as a recurrent blood loss, the underlying cause or causes should be corrected if possible.

As with all oral iron preparations, Ferrex™ 150 should be stored out of the reach of children to protect against accidental iron poisoning. Patients should not exceed the recommended dosage unless directed by a physician. Patients should be informed that iron products can cause dark or black stools.

Since oral iron products interfere with absorption of certain antibiotics, these products should not be taken within two hours of each other.

If pregnant, or planning to become pregnant or are currently breast-feeding please contact your physician, or health-care provider before using or continuing use.

51991-203-11

Ferrex™ 150

Polysaccharide-Iron Complex Capsules

**DO NOT USE IF BLISTER
SEALS ARE BROKEN**

breckenridge | A Towia
Company

Dietary Supplement
100 Unit Dose Capsules

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

ADVERSE REACTIONS

Adverse reactions with iron therapy may include constipation, diarrhea, nausea, vomiting, dark stool and abdominal pain. Adverse reactions with iron therapy are usually transient.

OVERDOSAGE

The clinical cause of acute iron overdose can be variable. Initial symptoms may include abdominal pain, nausea, vomiting, diarrhea, tarry stools, melena, hematemesis, hypotension, tachycardia, metabolic acidosis, hyperglycemia, dehydration, drowsiness, pallor, cyanosis, lassitude, seizures, shock and coma.

DOSAGE AND ADMINISTRATION

Usual adult dosage is one or two capsules daily, or as directed by a physician or health-care provider. **Do not exceed recommended dosage.**

HOW SUPPLIED

Ferrex™ 150 Capsules are supplied as orange/brown capsules, imprinted "B203". Each blister contains 10 capsules per card, 51991-203-11.

WARNING: Keep this and all medications out of the reach of children.

Store at 25°C (77°F); excursions permitted to 15° - 30°C (59° - 86°F). See USP Controlled Room Temperature. Protect from light and moisture.

These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

All substitutions using this product shall be pursuant to state statutes as applicable. This is not an Orange Book product.

To report a serious adverse event contact: 1-800-367-3395.

Manufactured by:

Contract Pharmacal Corp.
Hauppauge, NY 11788

Distributed by:

Breckenridge Pharmaceutical, Inc.
Berlin, CT 06037

⚠ WARNING: Reproductive Harm - www.P65Warnings.ca.gov.

11749-05-20

