

PE SCIENCE

4

Week Supply

500mg

Capsules per

10

Capsules per

# Mass-NP

## BULKING SUPPORT

### Formulated From Science

- Designed for Bulking Support
- Cinnulin PF<sup>®</sup>
- 4-OHile<sup>™</sup>  
4-Hydroxyisoleucine



MS-100000

DIETARY SUPPLEMENT

84 CAPSULES

## Supplement Facts

Serving Size: 3 Capsules  
Servings Per Container: 28

### Amount Per Serving %DV

Cinnulin PF <sup>®</sup> <small>(whole whole extract of cinnamon bark) Cinnamomum cassia with Cinnamomum lanceolatum (standardized to 7% Total Polyphenols)</small>	500 mg	*
Proprietary Blend <small>(standardized from bitter orange peel extract)</small>	250 mg	*
4-Hydroxyisoleucine <small>(standardized from Krugiodendron extract)</small>	100 mg	*
Ascorbic Acid <small>(standardized from lemon fruit extract)</small>	10 mg	*

\*Daily Value Not Established

Other Ingredients: Gelatin, Silicon Dioxide, Magnesium Stearate, FD&C Blue #1, FD&C Red #40, Titanium Dioxide  
Manufactured by PE Science LLC (3880 East Bay Dr #200-105, Lely, FL 32771, USA) (Ph: 888-885-0199)



0.1

Cinnulin PF<sup>®</sup> is a registered trademark of PE Science LLC

Directions: Take 3 capsules daily. For best results take 1 capsule, 3 times per day with a meal. Do not exceed 3 capsules per day.

**WARNING:** This product is intended to be consumed by healthy adults 18 years of age or older. It is designed for individuals interested in fitness who generally do not have health problems. Pregnant or nursing women should not use this product. Consult with a licensed physician before using this product, especially if you are taking any prescription, over the counter medication, dietary supplement product, or if you have any preexisting medical condition including but not limited to: high or low blood pressure, high or low cholesterol, cardiac arrhythmia, stroke, heart, liver, kidney or thyroid disease, seizure disorder, psychiatric disease, osteoporosis, diabetes, difficulty urinating due to prostate enlargement or if you are taking a MAO-B inhibitor or any other medication. Discontinue use 2 weeks prior to surgery. Discontinue use and immediately consult your health care professional if you experience any adverse reaction to this product. Do not exceed recommended serving. Do not use if safety seal is broken or missing.

KEEP OUT OF REACH OF CHILDREN.

These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.