

LIVE FOR THE PUMP!®

NITRIX® PRODUCT HIGHLIGHTS:

NITRIX® is designed to support and enhance:

- Sustained Increases in Nitric Oxide (N.O.) Production*
- Muscle Fullness, Vascularity and Full-Body Pumps*
- Muscle Strength, Power, Endurance*
- Nutrient Partitioning*
- Removal of Fatigue-Associated Waste Products from Muscle Tissue*
- Recovery from Exercise*
- Blood Flow to Muscle Tissue*
- Muscle Carnosine Content*
- Resistance to Muscular Fatigue*
- Hydrogen Ion (H⁺) Buffering*
- Anaerobic Working Capacity of Muscle Tissue*

Stack with ATRO-PHEX®, CELLMASS®, N.O.-XPLODE™, AXIS-HT™, SYNTHA-6™, and TRUE-MASS® for maximum physique and performance impact.*

DESIGNED FOR: Healthy persons seeking to support or enhance lean muscle, physical performance, and recovery.*

RECOMMENDED USE FOR PERSONS UNDER 200 LBS: As a dietary supplement, take 3 tablets 3 times per day, every day. Always take NITRIX® on an empty stomach (i.e. approximately 30-45 minutes before meals or 2 hours after a meal). Taking NITRIX® with food or on a full stomach may diminish its effects. Take the first 3 tablets in the morning upon rising, the second 3 tablets before lunch, and the third 3 tablets before dinner.

RECOMMENDED USE FOR PERSONS OVER 200 LBS: Follow the same guidelines as above, but take 4 tablets 3 times per day, every day, on an empty stomach.

NOTE: Do not take glutamine supplements and NITRIX® at the same time as doing so may diminish its effects. Keep intake of glutamine supplements to 10g or less of glutamine daily. For more info see Q&A section of www.BSNOnline.net. To maximize the vaso-muscular volumizing effects take NITRIX® 3 times per day, every day.* Also consume 120 oz. of water per day and a diet that is rich in protein and carbohydrates. Your body chemistry and weight will determine how long it takes to experience the effects of NITRIX®. Many users may notice the effects within the first 7-14 days of recommended use. Each week thereafter, the effects will become more pronounced. Other users may notice the initial effects after 14-21 days of use. Again, the effects will become more pronounced each week thereafter. NITRIX® begins to induce its maximum vaso-muscular volumizing effects in weeks 3-12.* After 12 weeks of use, discontinue for at least 4 weeks. Store in a cool dry place.

NITRIX® supports the following segments of the BSN Supplement Pyramid™: Muscle Support, Performance Support and Recovery Support.* For more information visit www.BSNOnline.net.



CORE SERIES



FINISH FIRST.®



NITRIX®

— A.M. TO P.M. VASO-MUSCULAR VOLUMIZER* —



SUPPLEMENT FACTS

Serving Size: 3 Tablets
Servings Per Container: 120

	Amount Per Serving	% DV
Folate (Folic Acid)	200mcg	50%
ACE3 (Triple Action Nitric Oxide Matrix) L-Arginine AKG, L-Arginine Ethyl Ester, L-Citrulline	3.0g	**
CRTS2 (Controlled Release Technology & Support System)* Hydroxypropyl Methylcellulose (A Micro-Polymer Hydrophilic Matrix)	1.5g	**
AVPT (Advanced Volumizing & Performance Technology)* Creatine Ethyl Ester-Beta-Alanine Composite (CarnoSyn®), Sodium Creatine Phosphate Matrix, Creatinol-O-Phosphate-Malic Acid Interfusion, Creatine AAB™ (Creatine-Alpha-Aminobutyric Acid Matrix)		**
Phosphoplexx Di-Calcium Phosphate, Di-Potassium Phosphate, & Sodium Phosphate		**
NAD (Nicotinamide Adenine Dinucleotide)		**

**Daily Value (DV) Not Established.

Other Ingredients: Magnesium Stearate and Stearic Acid.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

WARNING: Before consuming NITRIX® seek advice from a physician if you are unaware of your current health condition or have any pre-existing medical condition including but not limited to: high or low blood pressure, cardiac arrhythmia, stroke, heart, liver, or thyroid disease, anxiety, depression, seizure disorder, psychiatric disease, diabetes, pernicious anemia, difficulty urinating due to prostate enlargement, or if you are taking an MAO inhibitor or any other medication. Do not use if you are contemplating pregnancy, pregnant, nursing, prone to dehydration, or exposed to excessive heat. Reduce or discontinue use if sleeplessness, tremors, dizziness, nervousness, headaches, or heart palpitations occur. Combining NITRIX® with alcohol may increase the risk of low blood pressure (hypotension) and/or low blood sugar (hypoglycemia) particularly in the fasted state. Please be aware this product contains the naturally occurring amino acid Beta-Alanine, which may cause a tingling skin sensation in some individuals similar to niacin flush. This effect should diminish after a few hours and should ultimately subside after days of continuous use. Although rare, individuals with the genetic disorder hyper Beta-Alaninemia should not use this product. NITRIX® is only intended to be consumed by healthy adults 18-50 years of age. Keep out of reach of children and pets.

Allergen Warning: Manufactured on equipment which processes products containing milk, egg, soybeans, shellfish, fish, tree nuts and peanuts.

Natural Alternatives International (NAI) is the owner of patents 5,965,596; 6,172,098; 6,426,361; 6,680,294 and registered trademark CarnoSyn®.

Free Supplement Advice Hotline!
877.673.3727
www.BSNOnline.net

Developed By & Manufactured Exclusively For:
Bio-Engineered Supplements & Nutrition
Boca Raton, FL 33487 USA

WWW.BSNONLINE.NET

DIETARY SUPPLEMENT • 360 TABLETS

WWW.BSNONLINE.NET