

Becozinc



Dr.Reddy's



Each capsule contains:

Zinc Sulfate Monohydrate USP - 54.93 mg, Thiamine Nitrate BP - 10 mg, Riboflavin BP - 10 mg, Pyridoxine Hydrochloride BP 3 mg, Vitamin B12 coated, equivalent to Cyanobalamin Bp -15 mg, Ascorbic Acid BP -150 mg, Folic Acid Bp -1 mg, Nicotinamide BP - 50 mg Calcium Pantothenate BP - 12.5 mg

Storage

Store below 25° C, Protect from light and moisture*. Keep out of reach of children.

Precautions

Before using Becozinc Capsule, inform your doctor about your current list of medications, over the counter products (e.g. vitamins, herbal supplements, etc.), allergies, pre-existing diseases, and current health conditions (e.g. pregnancy, upcoming surgery, etc.). Some health conditions may make you more susceptible to the side-effects of the drug. Take as directed by your doctor or follow the direction printed on the product insert. Dosage is based on your condition. Tell your doctor if your condition persists or worsens. Important counseling points are listed below.

- Administration of folic acid alone is improper therapy for pernicious anemia and other megaloblastic anemias in which vitamin B12 is deficient
- Avoid alcohol consumption, especially if having alcoholic cirrhosis
- Avoid alcohol intake while treating with niacin
- Avoid taking in empty stomach
- Do not use if suffering from stomach ulcer
- Do not use if you are planning to be pregnant
- Drink plenty of liquids

Interactions with Becozinc Capsule

If you use other drugs or over the counter products at the same time, the effects of Becozinc Capsule may change. This may increase your risk for side-effects or cause your drug not to work properly. Tell your doctor about all the drugs, vitamins, and herbal supplements you are using, so that your doctor can help you prevent or manage drug interactions.

When not to use Becozinc Capsule

Hypersensitivity to Becozinc Capsule is a contraindication. In addition, Becozinc Capsule should not be used if you have the following conditions:

- Allergic reactions
- Breastfeeding
- Hypersensitivity
- Intolerance to the drug
- Leber's disease
- Pregnant
- Severe renal impairment

Composition and Active Ingredients

Becozinc Capsule is composed of the following active ingredients (salts)

- Folic Acid - Vitamin B9 - 1 MG
- Vitamin B12 - 15 MCG
- Zinc Sulphate - 54.93 MG
- Vitamin B1 - 10 MG
- Niacinamide - 50 MG
- Vitamin B2 - 10 MG
- Vitamin B6 - 3 MG
- Calcium Pantothenate - 12.5 MG

Overdosage of Becozinc Capsule

- Do not use more than prescribed dose. Taking more medication will not improve your symptoms; rather they may cause poisoning or serious side-effects. If you suspect that you or anyone else who may have overdosed of Becozinc Capsule, please go to the emergency department of the closest hospital or nursing home. Bring a medicine box, container, or label with you to help doctors with necessary information.
- Do not give your medicines to other people even if you know that they have the same condition or it seems that they may have similar conditions. This may lead to overdosage.
- Please consult your physician or pharmacist or product package for more information.

Storage of Becozinc Capsule

- Store medicines at room temperature, away from heat and direct light. Do not freeze medicines unless required by package insert. Keep medicines away from children and pets.
- Do not flush medications down the toilet or pour them into drainage unless instructed to do so. Medication discarded in this manner may contaminate the environment. Please consult your pharmacist or doctor for more details on how to safely discard Becozinc Capsule.

Dosage Information

Please consult your physician or pharmacist or refer to product package.

CONTACT

Phone: +95 1 557 477
+95 1 860 3825

Email: supportmyanmar@drreddys.com
Dr.Reddy's Laboratories Ltd.
Room No. (F1) to (F5),
First Floor, Block (A),
Pearl condominium,
Kabaraye Pagoda Road, Yangon.