

KEEP OUT OF
REACH OF
CHILDREN

NOT
RECOMMENDED
FOR USE BY
MINORS

Lot# 59-2798

DIET & ENERGY SUPPLEMENT

STACKER3

XPLC extreme
Performance
Formula

HELPS...

- BURN FAT[†]**
- INCREASE ENERGY[†]**
- BOOST METABOLISM[†]**

Stacker 3 XPLC combined with a low calorie, healthy diet and regular exercise may help you to achieve your desired goals.

4 Capsules



STACKER3
XPLC extreme
Performance
Formula

SUPPLEMENT FACTS		
SERVING SIZE: 1 CAPSULE	SERVINGS PER CONTAINER: 4	
	AMOUNT PER SERVING	% DV
CHROMIUM (AS CHROMIUM PICOLINATE)	200mcg	170%*
PROPRIETARY BLEND	235mg	**
YERBA MATE (LEAVES), CAMELLIA SINENSIS 50% POLYPHENOLS (GREEN TEA LEAVES), TAURINE, YOHIMBINE ALKALOIDS (PAUSINYSTALIA YOHIMBE BARK).		
CAFFEINE	200mg	**

* PERCENT DAILY VALUES (%DV) ARE BASED ON A 2,000 CALORIE DIET.
** DAILY VALUE NOT ESTABLISHED

OTHER INGREDIENTS: DEXTROSE, GELATIN, DIBASIC CALCIUM PHOSPHATE, STEARIC ACID, MAGNESIUM STEARATE, TITANIUM DIOXIDE, FD&C RED #3, BLUE #1, RED #40.

SUGGESTED USE: ONE CAPSULE AFTER MEALS. DO NOT EXCEED 3 CAPSULES IN A 24 HOUR PERIOD.

WARNING: NOT RECOMMENDED FOR USE BY MINORS. KEEP OUT OF REACH OF CHILDREN. DO NOT USE IF YOU ARE PREGNANT OR NURSING. CONSULT A PHYSICIAN OR LICENSED QUALIFIED HEALTH CARE PROFESSIONAL BEFORE USING THIS PRODUCT IF YOU HAVE, ARE AT RISK FOR, OR HAVE A FAMILY HISTORY OF STROKE, HEART DISEASE, THYROID DISEASE, LIVER DISEASE, KIDNEY DISEASE, ULCER, DIABETES, HIGH BLOOD PRESSURE, CAFFEINE SENSITIVITY, RECURRENT HEADACHES, ANXIETY, DEPRESSION OR OTHER PSYCHIATRIC CONDITION, GLAUCOMA, DIFFICULTY URINATING, PROSTATE ENLARGEMENT, SEIZURE DISORDER, OR IF YOU ARE USING ANY OTHER DIETARY SUPPLEMENT, OR A MONOAMINE OXIDASE INHIBITOR (MAOI), PRESCRIPTION DRUG OR OVER-THE-COUNTER DRUG CONTAINING EPHEDRINE OR PSEUDOEPHEDRINE (FOUND IN SOME ALLERGY ASTHMA, AND COUGH/COLD PRODUCTS). DO NOT EXCEED RECOMMENDED SERVING. INDIVIDUALS WHO EXCEED THE RECOMMENDED SERVING OR CONSUME CAFFEINE WITH THIS PRODUCT MAY EXPERIENCE SERIOUS ADVERSE HEALTH EFFECTS, INCLUDING MUSCLE FUNCTION LOSS, CHILLS AND VERTIGO. STOP USE AND CALL A PHYSICIAN OR LICENSED QUALIFIED HEALTH CARE PROFESSIONAL IMMEDIATELY IF YOU EXPERIENCE RAPID HEARTBEAT, NAUSEA, INSOMNIA, DIZZINESS, SEVERE HEADACHE, SHORTNESS OF BREATH, OR SIMILAR SYMPTOMS. IN CASE OF OVERDOSE SEEK PROFESSIONAL ASSISTANCE OR CONTACT A POISON CONTROL CENTER IMMEDIATELY. AVOID ALCOHOL AND FOODS WITH TYRAMINE (SUCH AS CHEESE, RED WINE AND LIVER) WHILE TAKING THIS PRODUCT.

MANUFACTURED & DISTRIBUTED BY NVE PHARMACEUTICALS
ANDOVER, NJ 07821 • TEL. 973-786-7868 • WWW.STACKER2.COM

[†] THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE
FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT
INTENDED TO DIAGNOSE, TREAT, CURE OR PREVENT DISEASE.

D121313



PART# 3765



8 78114 59001 0

IMPORTANT: READ INSTRUCTIONS ON REVERSE SIDE BEFORE USE.