

NDC 42937-705-10

Prenatal Plus Iron Tablets

Multivitamin/Multimineral
Supplement

For Use Before, During
and After Pregnancy.

Rx only

100 Tablets

Each Tablet Contains: VITAMINS:

Each Tablet Contains:	Amount per Tablet
Vitamin A (from Acetate and Beta Carotene)	4000 IU
Vitamin C (Ascorbic Acid)	120 mg
Vitamin D3 (Cholecalciferol)	400 IU
Vitamin E (d-Alpha Tocopheryl Acetate)	22 IU
Thiamine (Vitamin B-1 from Thiamine Mononitrate)	1.84 mg
Riboflavin (Vitamin B-2)	3 mg
Niacin (as Nicotinamide)	20 mg
Vitamin B6 (from Pyridoxine HCl)	10 mg
Folic Acid	1 mg
Vitamin B-12 (cyanocobalamin)	12 mcg

MINERALS:

Calcium (from Calcium Carbonate)	200 mg
Iron (from Carbonyl Iron)	29 mg
Zinc (from Zinc Oxide)	25 mg
Copper (from Cupric Oxide)	2 mg

Other Ingredients: Ascorbyl Palmitate, Citric Acid anhydrous, DL- alpha Tocopherol, Ethylcellulose, FD&C Blue #2 Lake, FD&C Red #40 Lake, FD&C Yellow #5 Lake, FD&C Yellow #6 Lake, Glucose, Gum Acacia, Hypromellose, Magnesium Stearate, Mize Starch, Maltodextrin, Methylcellulose, Microcrystalline Cellulose, Mineral Oil, Mono- and di-glycerides, Polyethylene Glycol, Pregelatinized Corn Starch, Silicon Dioxide, Sorbit Acid, Soy Protein, Stearic Acid, Sucrose, Titanium Dioxide, Tri calcium Phosphate.

Indications: To provide Vitamin and Mineral supplementation throughout pregnancy and during the postnatal period for both the lactating and nonlactating mother. It is also useful for improving nutritional status prior to conception.

Directions: As a dietary adjunct before, during and after pregnancy, take one tablet daily with a meal, or as directed by physician.

WARNING: A accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under age 6. Keep this product out of the reach of children. In case of accidental overdose, call a doctor or Poison Control Center immediately.

Precautions: Folic acid may partially correct the hematologic damage due to Vitamin B12 deficiency of pernicious anemia, while the associated neurological damage progresses.

This product contains FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No. 5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin type sensitivity.

Store at room temperature. Dispense in a well closed light-resistant container, with a child resistant cap.

CALL YOUR DOCTOR FOR MEDICAL ADVICE ABOUT SIDE EFFECTS. YOU MAY REPORT SIDE EFFECTS TO THE FDA AT 1-800-FDA-1088 (TOLL FREE).

Keep this and all medication out of the reach of children.

Do not use if imprinted Safety Seal under cap is broken or missing.

Distributed by: Nationwide Laboratories, Iselin NJ 08830

Lot No.: Exp. Date: Rev. 7/12

