

Supplement Facts

Serving Size: 1 Caplet
Servings Per Box: 100

	Amount per Serving	% DV for Pregnant and Lactating Women
Vitamin A (as Beta Carotene)	630 mcg RAE (100 IU)	45%
Vitamin C (Ascorbic Acid)	120 mg	100%
Vitamin D (as Cholecalciferol)	8 mcg (32 IU)	53%
Vitamin E (as Alpha-Tocopheryl Succinate)	13 mg* (20 IU)	64%
Thiamin (as Thiamine Mononitrate)	2 mg	163%
Riboflavin	3.4 mg	203%
Niacin (as Nicotinamide)	16 mg NE	54%
Vitamin B₆ (as Pyridoxine HCl)	10 mg	500%
Folate	2083 mcg DFE (1050 mcg Folate Acid)	147%
Vitamin B₁₂ (as Cyanocobalamin)	15 mcg	336%
Pantoic Acid	5 mg	71%
Iron (as Ferrous* Carbonate)	50 mg	183%
Magnesium (as Magnesium Oxide)	15 mg	4%
Zinc (as Zinc Oxide)	10 mg	39%
Copper (as Copper Gluconate)	1 mg	20%

* Equal to mcg alpha-tocopherol

Other Ingredients: Microcrystalline Cellulose, Di-Calc Phosphate, Croscarmellose Sodium, Pink Color Coating (Hydroxypropylmethyl Cellulose, Polyvinyl Alcohol, Titanium Dioxide, Polyethylene Glycol, Ink, FD&C Red #40 Lake), Fumed Silica, Stearic Acid, Magnesium Stearate, Povidone K30, Acacia.

GLUTEN, SUGAR, AND LACTOSE-FREE

Elite OB™ is a prescription multivitamin/multimineral indicated for use in improving the nutritional status of women prior to conception, throughout pregnancy, and in the postnatal period for both lactating and non-lactating mothers.

CONTRAINDICATIONS

Elite OB™ should not be used by patients with a known hypersensitivity to any of the listed ingredients.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. **KEEP THIS PRODUCT OUT OF REACH OF CHILDREN.** In case of accidental overdose, call a doctor or poison control center immediately.

PRECAUTIONS

Folate acid alone is an improper therapy in the treatment of pyridoxine deficiency and other megaloblastic anemias where vitamin B₁₂ is deficient. Folate acid in doses above 0.1 mg daily may obscure pyridoxine deficiency, in that hematologic remission can occur while neurological manifestations remain progressive. Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

The patient's medical conditions and consumption of other drugs, herbs, and/or supplements should be considered.

DRUG INTERACTIONS

Elite OB™ caplets are not recommended for and should not be given to patients receiving levodopa because the action of levodopa is antagonized by pyridoxine. There is a possibility of increased bleeding due to pyridoxine interaction with anticoagulants (e.g., Aspirin, Heparin, Clopidogrel).

ADVERSE REACTIONS

Adverse reactions have been reported with specific vitamins and minerals but generally at levels substantially higher than those contained herein. However, allergic and idiosyncratic reactions are possible at lower levels. Iron, even at the usual recommended level, has been associated with gastrointestinal intolerance in some patients.

DESCRIPTION

Elite OB™ is a pink capsule shaped caplet imprinted with "VP 010".

DIRECTIONS FOR USE

Take one caplet by mouth daily or as directed by a physician.

HOW SUPPLIED

Elite OB™ is available in a unit dose pack of 100 (10x10) caplets.

Product Code: 13811-027-10

STORAGE: Store at controlled room temperature 15°-30°C (59°-86°F). [See USP]. Protect from light and moisture and avoid excessive heat.

KEEP THIS PRODUCT OUT OF REACH OF CHILDREN.

For use on the order of a healthcare practitioner.

Call your doctor about side effects. To report side effects, call Trigen Laboratories, LLC at 1-888-8-TRIGEN (1-888-887-4436) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Customer Service: 1-888-887-4436 Rev. 06/2008

Manufactured for:

Trigen Laboratories, LLC
Bridgeport, NJ 08807
www.trigenlab.com



13811-027-10



Elite OB™ is a prescription multivitamin/multimineral indicated for use in improving the nutritional status of women prior to conception, throughout pregnancy, and in the postnatal period for both lactating and nonlactating mothers.



100 CAPLETS

13811-027-10



Elite OB™ is a prescription multivitamin/multimineral indicated for use in improving the nutritional status of women prior to conception, throughout pregnancy, and in the postnatal period for both lactating and nonlactating mothers.

