



PRENATAL PLUS IRON
Blenheim Pharmacal, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

Prenatal Plus Iron Tablets

DESCRIPTION

Prenatal Plus Iron Tablet is caplet shaped, film coated tablet debossed CIS 28, and provides 10 vitamins and 4 minerals to supplement the diet before, during and after pregnancy.

Each Prenatal Plus tablet Contains:

- Vitamin A (Acetate and Beta Carotene).....4000 I.U.
- Vitamin C (Ascorbic acid).....120 mg
- Vitamin D-3 (Cholecalciferol).....400 I.U.
- Vitamin E (dl- Alpha Tocopheryl Acetate).....22 I.U.
- Thiamine (Vitamin B-1 from Thiamine Mononitrate.....1.84 mg
- Riboflavin (Vitamin B-2).....3 mg
- Niacin(Niacinamide).....20 mg
- Vitamin B-6 (Pyridoxine HCl).....10 mg
- Folic Acid.....1 mg
- Vitamin B-12 (Cyanocobalamin).....12 mcg
- Calcium (Calcium Carbonate).....200 mg
- Iron (Carbonyl Iron).....29 mg
- Zinc (Zinc Oxide).....25 mg
- Copper (Cupric Oxide).....2 mg

Other ingredients: Ascorbyl Palmitate, Citric Acid anhydrous, DL-alpha Tocopherol, Ethylcellulose, FD&C Blue #2 lake, FD&C Red #40 lake, FD&C Yellow # 5 lake, FD&C Yello # 6 lake, Glucose, Gum Acacia, Hypromellose, Magnesium Stearate, Maize Starch, Maltodextrin, Methylcellulose, Microcrystalline Cellulose, Mineral Oil, Mono-and di-glycerides, Polyethyelen Glycol, Pregelatinized Corn Starch, Silicon Dioxide, Sorbic Acid, Soy Protein, Stearic Acid, Sucrose, Titanium Dioxide, Tricalcium Phosphate

INDICATIONS

To provide Vitamin and Mineral supplementation throughout pregnancy and during the postnatal period for both the lactating and non lactating mother. It is also useful for improving nutritional status prior to conception

DOSAGE

As a dietary adjunct before, during and after pregnancy, take one tablet daily with a meal, or as directed by a physician.

WARNING

Warning: Accidental overdose of iron containing products is a leading cause of fatal poisoning in children under age 6. Keep this product out of the reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

Precautions: Folic acid may partially correct the hematological damage due to Vitamin B12 deficiency of pernicious anemia, while the associated neurological damage progresses.

This product contains FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yello No. 5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

STORAGE

Store at room temperature. Dispense in a well closed light resistant container with a child resistant cap.

HOW SUPPLIED

NDC 0904-5339-60 plastic bottle containing 100 Tablets.

Principal Display Panel

Prenatal Plus Iron Tablets (Multivitamin/Multimineral Supplement)
31 Tabets
10544-609-31