Multivitamin/Mineral Softgel

Essential Nutrition for the Three Stages of Pregnancy: Preconception • Pregnancy • Postnatal

Rx Only

NDC 67112-402-30

30 Softgels

ONCE DAILY

DHA ONE EX

TriCare Prenatal DHAone.com

500 mg 215 mg

45 mg

3 mg 3.4 mg

20 mg

25 mg

100 mcg

27 mg 10 mg 2 mg 150 mcg

Description: Each purple softgel contains: Essential Fatty Acids Fish Oil Concentrate 10/50 TG

ti Oil Concenti DHA EPA

Vitamins
Vitamin B-1 (Thiamine Mononitrate)

Vitamin B-1 (Thiamine Monoritrate)
Vitamin B-2 (Riboflavin)
Vitamin B-3 (Inositol Hexanicctinate)
Vitamin B-6 (Pyridoxine Hydrochloride)

Vitamin B-12 (Cyanocobalamin) Folic Acid

Folic Acid
Vitamin C (Ascorbic Acid)
Vitamin D3 (Cholecalciferol in Corn Oil)
Vitamin E (d-alpha-Tocopherol)

Biotin

Iron (Ferrous Furnarate) Zinc (Zinc Sulfate Monohydrate) Copper (Cupric Sulfate Anhydrous)

lodine (Potassium lodide)
Other Active Ingredients

Docusate Sodium (with 15% sodium benzoate) 25 mg

INACTIVE INGREDIENTS: Gelatin Capsule (Bovine Gelatin, Glycerin, Sorbitol
Sorbitan Solution, Purified Water, Titanium Dioxide, FD&C Red #40, FD&C

Red #3, FDGC Blue #1), Yellow Beeswax, Soy Lecithin
INDICATIONS AND USAGE: TriCare Prenatal™ DHA ONE® RX

Multivitamin/Mineral Softgels are indicated to provide multivitamin/mineral/essential fatty acid supplimentation throughout pregnancy and during the postnatal period for both the lactating and non-lactating mother. It is also useful for improving nutritional status prior to conception.

CONTRAINDICATIONS: This product is contraindicated in patients with a known hypersensitivity to any of the ingredients.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctar or poison control center immediately.

Folic acid alone is improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where vitamin B-12 is deficient.

NOTICE: Contact with moisture may produce surface discoloration or erosion of the softgel.

PRECAUTIONS: Folic acid in doses above 10 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations progress.

ADVERSE REACTIONS: Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

DOSAGE AND ADMINISTRATION: One softgel daily or as directed by

A DISTANCE OF THE PROPERTY OF

(NDC 67112-402-30)

to bleeding.

WARNING: logestion of more than 3 grams of Omega 3 fatty actifs (such as DHA) per day has been shown to have potential, annihirombotic effects, including an increased bleeding time and international Normalized Ratio (NR). Administration of Omega 3 fatty actifs should be avoided in patients taking anticoagulants and in those known to have inherited or acquired predisposition.

Store at controlled room temperature of: 20°-30°C (68°-77°F). Excursions permitted to 15°-30°C (59°-86°F). KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

Marketed by Medecor Pharma LLC. Baton Rouge, LA



