



A **small** yet **mighty™**
once-daily prenatal vitamin

Nutritional Information

Prenate Pixie® – An innovative approach in prenatal nutrition.

- A small yet mighty™ once-daily softgel
- Sized and designed for preconception
- Packed with robust nutrients to carry patients through every step of pregnancy

VITAMINS	
Folate (as (6S)-N5-methyltetrahydrofolic acid calcium salt (equivalent to 600 mcg of folic acid) and folic acid, USP 400 mcg)†	1700 mg DFE
Vitamin B6 (as pyridoxine HCl)	5 mg
Vitamin B12 (as cyanocobalamin)	13 mcg
Biotin	75 mcg
Vitamin D3 (as cholecalciferol)	12.5 mcg
Vitamin C (as ascorbic acid)	30 mg
Vitamin E (as dl-alpha tocopheryl acetate)	4.5 mg
MINERALS	
Iron (as Sumalate® (ferrous asparto glycinate))	10 mg
Iodine (as potassium iodide)	150 mcg
OTHER NUTRIENTS	
Docosahexaenoic acid (DHA)	200 mg
Blueberry extract (vaccinium spp.)	5 mg

THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION.
THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.

†Prenate Pixie® contains L-methylfolate calcium. The L-methylfolate ((6S)-N5-methyltetrahydrofolic acid calcium salt) in Prenate Pixie® is less than 1.0% D-isomer.

Sumalate® is a registered trademark of Albion Laboratories, Inc., covered by one or more claims of U.S. Patent Nos. 6,716,814; 8,007,846; and 8,425,956.

PRENATE®, Prenate Mini®, Prenate Pixie®, Prenate® Enhance, Prenate® Restore, Prenate Chewable®, Prenate DHA®, Prenate Elite®, and Prenate Essential® are registered trademarks of Avion Pharmaceuticals, LLC. All rights reserved.

Avion PHARMACEUTICALS
Marketed by Avion Pharmaceuticals, LLC, Alpharetta, GA 30005
1.888.61.AVION (1.888.612.8466) © 2021 Avion Pharmaceuticals

IMPORTANT SAFETY INFORMATION

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

WARNING: Ingestion of more than 3 grams of omega-3 fatty acids (such as DHA) per day has been shown to have potential antithrombotic effects, including an increased bleeding time and International Normalized Ratio (INR). Administration of omega-3 fatty acids should be avoided in patients taking anticoagulants and in those known to have an inherited or acquired predisposition to bleeding.