
TestroVax

[testrothione[™]] 2700 m

A group of 43 men were divided into two groups: 1) the functional TestroVax compound group and 2) an indistinguishable placebo group. All subjects consumed their test product every morning at breakfast for 12 consecutive days, and blood samples were taken before treatment, after 6 and 12 days of treatment, and 3 days after stopping treatment. All blood samples were drawn between 9:30-10:30 a.m., a time of day when serum oscillations of testosterone are at their minimum. Results showed that serum testosterone levels in the experimental group were significantly increased compared to those of the placebo group over the course of the study, wherein the key TestroVax compound led

to a 42% increase in testosterone after 12 days of use. Interestingly, 3 days after the suspension of treatment, testosterone was still increased 1.22 fold compared to baseline in the treatment group, showing a lasting effect.*



NOTE: For additional information on this or any other Novex Biotech product, please call our experienced customer service representatives at: 1-800-898-5153.

Store at controlled room temperature: 15° to 30°C / 59° to 86°F.

Product resale allowed only through authorized representatives. International distribution is not authorized without prior written approval.

Individual results will vary.

Our 100% satisfaction guarantee:

Because no product or program will work for everyone, TestroVax™ is backed by our 100% money-back guarantee. For details regarding our guarantee, please visit www.NovexBiotech.com or call customer service at: 1-800-898-5153.



Manufactured in the USA.

NOTICE: This product should not be consumed for more than 4 consecutive weeks with a 2 to 4 week washout period between cycles.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

©2012 ALL RIGHTS RESERVED BR14082-



TestroVax

[testrothione™] 2700 mg

CLINICALLY TESTED
TESTOSTERONE
BOOSTING COMPOUND*

42 1%
Increase in Serum
Testosterone Levels
In Just 12 Days!

HEALTHY TESTOSTERONE LEVELS ASSOCIATED WITH:

- MUSCLE MASS GROWTH
- INCREASED ENERGY
- INTENSIFIED LIBIDO



DIETARY SUPPLEMENT

90 Tablets

Supplement Facts

Serving Size: 3 Tablets Servings Per Container: 30

| | Amount Per Serving | % Daily Value‡ |
|---|-----------------------|-------------------|
| Vitamin B6 (as pyridoxine HCI) | 2 mg | 100% |
| Folic Acid | 400 mcg | 100% |
| Vitamin B12 (as cyanocobalamin) | 6 mcg | 100% |
| Testrothione", A Proprietary Blend Containing: | 2.7 g | |
| (D)-Asparaginic Acid | | t |
| Fenugreek (<i>Trigonella foenum-graecum</i>) (seed Powder |) | † |
| (S)-2-Amino-5-guanidinopentanoic Acid HCl | | t |
| Resveratrol (Polygonum cuspidatum) (root) Ext | ract | t |
| Grape (seed) Extract | | † |
| Grape (skin) Extract | | † |
| Rhodiola rosea (root) Extract | | † |
| Licorice (Glycyrrhiza glabra) (root) Powder | | † |
| Red Clover (blossoms) Powder | | t |
| Damiana (leaf) Extract | | † |
| Withania somnifera (root) Extract | | t |
| 2-Aminoethanesulfonic Acid | | † |
| Schizonepeta (aerial parts) Powder | | t |

‡ Percent Daily Values are based on a 2,000 calorie diet † Daily Value not established

Other Ingredients May Include: Microcrystalline Cellulose.

Hydroxypropylmethylcellulose, Stearic Acid, Croscarmellose Sodium, Silicon Dioxide, Magnesium Stearate, Polyvinyl Alcohol, Talc, Polyethylene Glycol, Titanium Dioxide, Polysorbate 80, Blue #2.

DIRECTIONS FOR USE: Take 3 tablets daily with breakfast.

WARNING: Do not exceed suggested daily serving. For adult males only. Before consuming TestroVax, consult your physician. Do not use TestroVax if you have any pre-existing medical condition, including but not limited to: high or low blood pressure, cardiac arrhythmia, stroke, heart, liver, or thyroid disease, anxiety, depression, seizure disorder, psychiatric disease, diabetes, pernicious anemia, difficulty urinating due to prostate enlargement, or if you are taking an MAO inhibitor or any other medication. Discontinue use if sleeplessness, tremors, dizziness, nervousness, headache, or heart palpitations occur. Keep out of reach of children. Use only as directed.

CAUTION: Increased testosterone levels may increase aggression, over-confidence, and sexual arousal.

Manufactured for Novex Biotech? LLC, Salt Lake City, UT 84116 If you or your health-care provider have any questions, please call: 1-800-898-5153.

Novex Biotech.