

60258-196-01

Prenatal 19

Tablets

Dietary Supplement

For Professional Use Only

CYPRRESS

PHARMACEUTICAL, INC.

100 Tablets

DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Supplement Facts

Serving Size: 1 Tablet

Amount per serving		%Daily Value**
Vitamin A (Beta Carotene)	1000 IU	13%
Vitamin C (Ascorbic Acid)	100 mg	167%
Vitamin D-3 (Cholecalciferol)	400 IU	100%
Vitamin E (dl-Alpha Tocopheryl Acetate)	30 IU	100%
Vitamin B-1 (Thiamine HCl)	3 mg	176%
Vitamin B-2 (Riboflavin)	3 mg	150%
Niacinamide (Vitamin B-3)	15 mg	75%
Vitamin B-6 (Pyridoxine HCl)	20 mg	800%
Folic Acid	1 mg	125%
Vitamin B-12 (Cyanocobalamin)	12 mcg	150%
Pantothenic Acid (Calcium Pantothenate)	7 mg	70%
Calcium (Calcium Carbonate)	200 mg	15%
Iron (Ferrous Fumarate)	29 mg	161%
Zinc (Zinc Oxide)	20 mg	133%
Docusate Sodium	25 mg	*

* Daily Value not established.

** Daily Value for Pregnant and Lactating Women.

Other Ingredients: Microcrystalline Cellulose, Croscarmellose Sodium, Maltodextrin, Hygromellose, Starch, Stearic Acid, Titanium Dioxide, Pharmaceutical Glaze, Dicalcium Phosphate, Acacia, Crospovidone, Sodium Lauryl Sulfate, Citric Acid, Magnesium Silicate, Magnesium Stearate, Triacetin, Sucrose, Soy Polysaccharide, Sodium Benzoate, Mineral Oil, Polyvinylpyrrolidone, Silica, Sodium Ascorbate, Mixed Tocopherols, Triglycerides and Tricalcium Phosphate. This product contains soy.

DIRECTIONS: One tablet daily or as directed by a physician.

CONTRAINDICATIONS: Prenatal 19 is contraindicated in patients with a known hypersensitivity to any of the components of this product. Hemochromatosis and hemosiderosis are contraindications to iron therapy.

WARNINGS:

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of the reach of children. In case of accidental overdose, call a doctor or Poison Control Center immediately.

Folic acid alone is improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where Vitamin B-12 is deficient.

PRECAUTIONS: Folic acid in doses above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations remain progressive. Pernicious anemia should be excluded before using this product since folic acid may mask the symptoms of pernicious anemia.

ADVERSE REACTIONS: Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

STORAGE: Store at room temperature, USP.

Keep out of the reach of children.

To report a serious adverse event or to obtain product information, contact (800) 856-4393.

Distributed By:
Cypress Pharmaceutical, Inc.
135 Industrial Blvd.
Madison, MS 39110

L480 Rev. 01/11



NON-VARNISH PLATE GOES HERE