

Modified Citrus Pectin Powder

Modified citrus pectin is prepared by a proprietary process, which reduces the molecular weight of the pectin molecule. Average molecular weight: 98kD. Methoxylation: not more than 5%.

Suggested Use

As a dietary supplement, 5 to 15 grams daily: dissolve one rounded teaspoon (approx. 5 g) of powder in water, juice or other liquid, one to three times daily, or as directed by a healthcare practitioner. Best taken at least 30 minutes before, or 1.5 hours after food.

CAUTION: Those on a sodium-restricted diet should consult their healthcare practitioner before use.

Variations in product color may occur. Keep in a cool, dry place, tightly capped.



Modified Citrus Pectin

Reduced Molecular Weight



dietary supplement

HYPOALLERGENIC

454 grams (16 oz.)

Supplement Facts

Serving Size 1 Teaspoon (5 g)
Servings Per Container 90

Amount Per Serving

Calories 20

% Daily Value*

Sodium	300 mg	13%
Potassium	10 mg	<1%
Total Carbohydrates	5 g	2%
Dietary Fiber	3.75 g	13.4%
Total Sugars	500 mg	10%
Added Sugars	0 mg	†
Modified Citrus Pectin	5 g	†
Total Polysaccharides	3.5 g	†
Galacturonic acid	2.75 g	†

* Percent Daily Values are based on a 2,000 calorie diet.

† Daily Value not established.

Formulated exclusively for Allergy Research Group®
South Salt Lake, UT 84115
www.allergyresearchgroup.com
Phone: 800.545.9960