

Modified Citrus Pectin Powder

Modified citrus pectin is prepared by a proprietary process, which reduces the molecular weight of the pectin molecule. Average molecular weight: 98kD. Methoxylation: not more than 5%.

Suggested Use

As a dietary supplement, 5 to 15 grams daily: dissolve one rounded teaspoon (approx. 5 g) of powder in water, juice or other liquid, one to three times daily, or as directed by a healthcare practitioner. Best taken at least 30 minutes before, or 1.5 hours after food.

CAUTION: Those on a sodium-restricted diet should consult their healthcare practitioner before use.

Variations in product color may occur. Keep in a cool, dry place, tightly capped.



Formulated exclusively
for NutriCology®
South Salt Lake, UT 84115
www.nutricology.com
Phone: 800.545.9960



Modified Citrus Pectin Powder

Reduced Molecular Weight

Hypoallergenic

Dietary Supplement

454 grams (16 oz.)

Supplement Facts

Serving Size 1 Teaspoon (5 g)
Servings Per Container 90

Amount Per Serving

Calories 20

% Daily Value*

Sodium 300 mg 13%

Potassium 10 mg <1%

Total Carbohydrates 5 g 2%

Dietary Fiber 3.75 g 13.4%

Total Sugars 500 mg 10%

Added Sugars 0 mg †

Modified Citrus Pectin 5 g †

Total Polysaccharides 3.5 g †

Galacturonic acid 2.75 g †

* Percent Daily Values are based on a 2,000 calorie diet.

† Daily Value not established.