



Actual Color and Size

## With Super Chelates Mith Super Chelates Mith Super Chelates

Suggested Use: Take two (2) tablets daily with food. Do not exceed recommended daily dose unless directed by your health care practitioner.

Keep out of reach of children in a cool, dry place.

## Supplement Facts

Serving Size = Two (2) tablets

Servings Per Container=Varied

|   | Amount Per Serving | % Daily Value |
|---|--------------------|---------------|
| \ru \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ | 4.5                | 20/           |
| Vitamin B12(Cyanocobalamin)             | 45 mcg             | 2%            |
| Calcium (Calcium Carbonate)             | 500 mg             | 38%           |
| Iron (Ferrous Fumarate)                 | 18 mg              | 100%          |
| Iodine (Potassium Iodide)               | 225 mcg            | 150%          |
| Magnesium (Magnesium Oxide)             | 250 mg             | 60%           |
| Zinc (Zinc Citrate)                     | 15 mg              | 136%          |
| Selenium (L-Selenomethionine)           | 70 mcg             | 127%          |
| Copper (Copper Gluconate)               | <u>2 mg</u>        | 222%          |
| Manganese (Manganese Citrate)           | 4 mg               | 174%          |
| Chromium (Chromium Polynicotii          |                    | 343%          |
| Molybdenum (Sodium Molybdate            | e) 75 mcg          | 167%          |
| Potassium (Potassium Carbonate,         | lodide) 45 mg      | <1%           |
|   |                    |               |
| Cobalt (Cyanocobalamin)                 | 2 mcg              | *             |
|   |                    |               |
| * Daily Value Not Established           |                    |               |

Other Ingredients: Cellulose, and stearic acid.

This product may be produced in a facility that also makes products containing milk, eggs, fish, crustacean shellfish, tree nuts, peanuts, and soy. Food supplements should not be used as a substitute for a varied diet.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of the reach of children. In case of accidental overdose, call a doctor or poison control



www.vitalbulk.com 294 Anna St., Watsonville, CA 95076 In case of adverse event: 855-885-2855