



MAJOR 

Prenatal Tablets

A Dietary Supplement For
Pregnant & Lactating Women



30 Tablets

Supplement Facts

Serving Size: 1 Tablet

Amount Per Serving		% Daily Value
Vitamin A (as Acetate and Beta Carotene)	4000 I.U.	80%
Vitamin C (as Ascorbic Acid)	100 mg.	167%
Vitamin D-3 (Cholecalciferol)	400 I.U.	100%
Vitamin E (as dl-Alpha Tocopheryl Acetate)	11 I.U.	37%
Vitamin B-1 (as Thiamine Mononitrate)	↓ 1.5 mg.	88%
Vitamin B-2 (Riboflavin)	1.7 mg.	85%
Niacinamide (as Vitamin B-3)	35 mg.	50%
Vitamin B-6 (as Pyridoxine HCl)	2.6 mg.	104%
Folic Acid	↓ 0.4 mg.	100%
Vitamin B-12 (as Cyanocobalamin)	4 mcg.	50%
Calcium (as Calcium Carbonate)	200 mg.	10%
Iron (as Ferrous Fumarate)	27 mg.	150%
Zinc (as Zinc Oxide)	25 mg.	167%

PULL BACK FOR MORE INFORMATION

WARNING: If you are pregnant, nursing, taking any medications or planning any medical procedure, consult your doctor before use. Discontinue use and consult your doctor if any adverse reactions occur.

KEEP OUT OF THE REACH OF CHILDREN. STORE IN A DRY PLACE AND AVOID EXCESSIVE HEAT. TAMPER RESISTANT. DO NOT USE IF SEAL UNDER CAP IS BROKEN OR MISSING.

Distributed by:
Major Pharmaceuticals
31775 Enterprise Drive
Livonia, MI 48150, USA
TO REPORT SIDE EFFECTS
CALL:
(800) 616-2471 M-29

Rev 02/10

OTHER INGREDIENTS: Calcium Sulfate, Microcrystalline Cellulose, Stearic Acid, Croscarmellose Sodium, Hydroxypropyl Cellulose, Acacia, Maltodextrin, Titanium Dioxide, Starch, Traces Magnesium Stearate, Magnesium Silicate, Dicalcium Phosphate, Mineral Oil, Sucrose, Silica, Sodium Acetate, Triethylsiloxane, FD&C Red #40 Lake, FD&C Yellow #6 Lake, Mixed Tocopherols, Sodium Benzoate, Sorbic Acid, Thioctic Acid, Thioctic Phosphate, BHT and Ascorbyl Palmitate.

DIRECTIONS: For adults, take one (1) tablet daily, preferably with a meal. As a reminder, discuss the supplements and medications you take with your health care provider.

FREE OF: Yeast, wheat, milk or milk derivatives, lactose, soy, artificial flavor, sodium less than 5 mg per serving.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under six. **KEEP THIS PRODUCT OUT OF THE REACH OF CHILDREN.** In case of accidental overdose, call a doctor or Poison Control Center immediately.