

UNDA **plex**
ANGIplex™

NDC 62106-1229-3

UNDA **plex**
ANGIplex™

Homeopathic Preparation

For the temporary relief of sore throat and dry throat.

30 Tablets

Seroyal UNDA
Made in Belgium by UNDA
Dist. by Seroyal USA
Pittsburgh, PA 15275
1-888-737-6925

V1.0

BARCODE LOCATION
883196 32740

Drug Facts

Active ingredients

Purpose

Each tablet contains:

Aconitum napellus (Aconite) Whole Plant 8X. Dry throat
Apis mellifica (Honey bee) 8X Difficulty swallowing
Arum triphyllum (Indian turnip) Underground Parts 4X . . . Sore throat
Belladonna (Deadly nightshade) Whole Plant 8X . . . Sore throat
($< 1.25 \times 10^{-10}$ % alkaloids, calculated)
Kalium bichromicum (Potassium bichromate) 10X. . . Sore throat
Lachesis mutus (Bushmaster) Venom 10X Throat pain
Mercurius solubilis (Hahnemann's soluble mercury) 10X. . . Sore throat
Phytolacca decandra (Pokeweed) Root 8X Sore throat
X is a homeopathic dilution: see www.seroyal.com/homeopathics for details.

Uses For the temporary relief of symptoms associated with
■ sore throat ■ dry throat

Product "Uses" are based on Homeopathic Materia Medica and have not been clinically tested. These "Uses" have not been evaluated by the Food and Drug Administration.

Warnings

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult physician promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by physician.

Stop use and ask a doctor if symptoms continue to persist after three days of use or are accompanied by a fever.

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

If pregnant or breastfeeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ Adults and adolescents (12 years and older): At the first sign of symptoms, dissolve one tablet under the tongue every half hour to one hour. ■ When symptoms subside, decrease frequency to every four hours. ■ Once the symptoms are gone, continue taking one tablet daily for three more days. ■ Children (under 12 years): Take under the direction of your healthcare practitioner.

Other information ■ Do not use if the blister pack has been tampered with. ■ Store in a cool, dry place.

Inactive ingredients Lactose monohydrate (milk), magnesium stearate, sodium carboxymethylcellulose, xylitol