

0.25 mg FLUORIDE
MULTIVITAMIN
LIQUID DROPS

NDC 52796-170-50

BonGeo.
PHARMACEUTICALS

Floriva
Plus Liquid Drops.
0.25 mg

Fluoride Multivitamin Liquid Drops

Quatrefolic[®]

Patent Refs: U.S. Patent No. 7,947,682;
Other patents pending



Natural Berry Flavor

Physician Prescribed
Dietary Supplement

1½ FL. OZ (50 mL)

NO ARTIFICIAL COLORS OR FLAVORS

SUGAR, SACCHARIN
AND GLUTEN FREE

*contains 0.25 mg fluoride per 1 mL from 0.55 mg sodium fluoride
Quatrefolic[®] Trademark and Patent property of Gnosis SpA.

Supplement Facts

Serving Size: 1 mL

Servings Per Container: 50

	Amount Per Serving	% Daily Value	
		1-12 months	1-4 years
Vitamin A (as retinyl palmitate)	1150 IU	77%	46%
Vitamin C (as ascorbic acid)	32 mg	91%	80%
Vitamin D3 (as cholecalciferol)	400 IU	100%	100%
Vitamin E (as d-alpha-tocopheryl succinate)	5 IU	100%	60%
Vitamin B1 (as thiamine HCl)	0.5 mg	100%	71%
Vitamin B2 (as riboflavin phosphate sodium)	0.8 mg	100%	75%
Niacin (as niacinamide)	2 mg	25%	22%
Vitamin B6 (as pyridoxine HCl)	0.4 mg	100%	57%
Folate (as 55 mcg Quatrefolic [®]) (5S-5-methyltetrahydrofolic acid, glucosamine salt)	29.7 mcg	30%	15%
Vitamin B12 (as cyanocobalamin)	2 mcg	100%	67%
Fluoride (as sodium fluoride)	0.25 mg	†	†

(†)Daily Value not established

Other Ingredients: Glycerin, purified water, propylene glycol, polysorbate 80, gum acacia, sucrose, maltodextrin, sodium hydroxide, potassium acetate, sodium benzoate, natural flavors.

WARNING: KEEP THIS PRODUCT OUT OF THE REACH OF CHILDREN. IF OVERDOSSAGE IS SUSPECTED, SEEK PROFESSIONAL ASSISTANCE OR CONTACT A POISON CONTROL CENTER IMMEDIATELY 1-800-222-1222.

Directions: Use as directed by a physician.
SHAKE WELL BEFORE USE.

How to Use: Fill dropper to appropriate line and dispense with a single squeeze of the dropper bulb. The full dose will be given. It is normal for a small amount to remain in the tip of the dropper.

Do not accept if bottle tamper seal is broken or missing.

Caution: Do not use this product if you are allergic to any of the ingredients. Take this product at least 2 hours before or after taking any products containing calcium (including milk, yogurt, other dairy products) or aluminum/magnesium hydroxide (e.g., certain antacids/acid-reducers) since these may decrease effectiveness. Prolonged daily ingestion of excessive fluoride may result in varying degrees of dental fluorosis.

OTHER INFORMATION: You may report serious side effects to 1-844-640-7607. BonGeo Pharmaceuticals, Inc. requires this product to be dispensed only under the order of a physician or licensed medical practitioner.

© 2016

BonGeo Pharmaceuticals, Inc.

101 Eisenhower Parkway #300, Roseland, NJ 07068

FPL02 03/13

