

DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Each Tablet Contains

Serving Size: 1 Tablet

Amount per serving	%Daily Value*
Vitamin A (Beta Carotene)	4000 IU 80%
Vitamin C (Ascorbic Acid)	120 mg 200%
Vitamin D-3 (Cholecalciferol)	400 IU 100%
Vitamin E (dl-Alpha Tocopheryl Acetate)	30 IU 100%
Vitamin B-1 (Thiamine Mononitrate)	3 mg 176%
Vitamin B-2 (Riboflavin)	3 mg 160%
Niacinamide (Vitamin B-3)	20 mg 100%
Vitamin B-6 (Pyridoxine HCl)	3 mg 120%
Folic Acid	1 mg 126%
Vitamin B-12 (Cyanocobalamin)	6 mcg 100%
Biotin	30 mcg 10%
Pantothenic Acid (Calcium Pantothenate)	7 mg 70%
Calcium (Calcium Carbonate)	200 mg 15%
Iron (Carbonyl Iron)	28 mg 181%
Iodine (Potassium Iodide)	150 mcg 100%
Magnesium (Magnesium Oxide)	100 mg 22%
Zinc (Zinc Oxide)	15 mg 100%
Copper (Cupric Oxide)	3 mg 150%

*Daily Value for Pregnant and Lactating Women.

OTHER INGREDIENTS:

Microcrystalline Cellulose, Sacchar 1500, Pink Coating (Polyethylene Glycol, Polyvinyl alcohol, Titanium Dioxide, Talc, FD&C Blue #1, FD&C Red #40 and FD&C Yellow #6), Colloidal Silicon Dioxide, Magnesium Stearate.

INDICATIONS: To provide vitamin and mineral supplementation throughout pregnancy and during the postnatal period for both the lactating and nonlactating mother.

DIRECTIONS: One tablet daily or as directed by a physician. Each tablet is pink colored, oval-shaped, and embossed with "RP204" on one side.

CONTRAINDICATIONS: This product is contraindicated in patients with a known hypersensitivity to any of the components of this product. Hemochromatosis and hemodialysis are contraindications to iron therapy.

WARNINGS:

Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of the reach of children. In case of accidental overdose, call a doctor or Poison Control Center immediately.

CAUTION: Folic acid alone is improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where Vitamin B-12 is deficient.

PRECAUTIONS: Folic acid in doses above 0.1 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations remain progressive. Pernicious anemia should be excluded before using this product since folic acid may mask the symptoms of pernicious anemia.

ADVERSE REACTIONS: Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

STORAGE: Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C, (between 59°F and 86°F).

KEEP OUT OF REACH OF CHILDREN.

To report a serious adverse event or to obtain product information, contact 1-888-438-1772.

Manufactured by
Rochester Pharmaceuticals
Rochester, NY 14624
www.RochesterPharm.com

834940 12/12

