

CORVITE[®]

150

Now with



TABLETS



Iron Supplementation

PATENT PENDING
Sugar, Lactose & Gluten Free
Vegan Safe

30 Tablets



DOSAGE & ADMINISTRATION
Usual adult dose is 1 tablet daily,
or as directed by a physician.

STORAGE

Store at controlled room temperature 15°-30°C (59°-86°F) [see USP]. Protect from light and moisture and avoid excessive heat. Dispense in a tight, light resistant container as defined by the USP using a child-resistant closure.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. **KEEP THIS PRODUCT OUT OF REACH OF CHILDREN.** In case of accidental overdose, call a doctor or poison control center immediately.



3 68025-052-30 6

Supplement Facts

Servings per bottle: 30

Serving Size: 1 tablet		% DV Adults & Children 4 Years or More
Vitamin C (calcium ascorbate)	120 mg	200%
Calcium Threonate	5 mg	0.5%
Vitamin B ₆ (pyridoxine HCl)	10 mg	500%
Vitamin B ₇ (biotin 1%)	100 mcg	33%
Vitamin B ₁₂ (cyanocobalamin)	20 mcg	333%
Folate	1 mg	250%
[as folic acid 700 mcg and Quatrefolic [®] ((6S)-5-methyltetrahydrofolic acid) 300 mcg, (equivalent to 600 mcg of (6S)-5-methyltetrahydrofolic acid, glucosamine salt)]**		
Iron (ferrous carbonyl iron powder)	150 mg	833%
Magnesium (oxide)	10 mg	3%
Zinc (citrate)	25 mg	167%

** Quatrefolic[®] is a registered trademark of Gnosis, SpA.
U.S. Patent No. 7,947,662.

Other Ingredients: Microcrystalline Cellulose, Pregelatinized Starch, Di-Calcium Phosphate, Red Color Coating (Hydroxypropylmethyl Cellulose, Polyvinyl Alcohol, FD&C Red #40 Lake, Polyethylene Glycol, Talc, Titanium Dioxide), Croscarmellose Sodium, Stearic Acid, Magnesium Stearate.

Sugar, Lactose, Gluten Free & Vegan Safe

CLINICAL PHARMACOLOGY: Iron is an essential component in the formation of hemoglobin. Adequate amounts of iron are necessary for effective erythropoiesis. Iron also serves as a cofactor of several essential enzymes, including cytochromes that are involved in electron transport. Folic acid is required for nucleoprotein synthesis and the maintenance of normal erythropoiesis. Folic acid is converted in the liver and plasma to its metabolically active form, tetrahydrofolic acid, by dihydrofolate reductase. Vitamin B₁₂ is required for the maintenance of normal erythropoiesis, nucleoprotein and myelin synthesis, cell reproduction and normal growth. Intrinsic factor, a glycoprotein secreted by the gastric mucosa, is required for active absorption of Vitamin B₁₂ from the gastrointestinal tract.

INDICATIONS AND USAGE: CORVITE[®]150 is indicated for the prevention and treatment of iron deficiency anemia and/or nutritional megaloblastic anemias.

CONTRAINDICATIONS: CORVITE[®]150 is contraindicated in patients with a known hypersensitivity to any of the components of this product. Hemochromatosis and hemosiderosis are contraindications to iron therapy.

WARNINGS: Folic acid alone is improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where Vitamin B₁₂ is deficient.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. **KEEP THIS PRODUCT OUT OF REACH OF CHILDREN.** In case of accidental overdose, call a doctor or poison control center immediately.

PRECAUTIONS

General: Do not exceed recommended dose. The type of anemia and the underlying cause or causes should be determined before starting therapy with CORVITE[®]150. Since the anemia may be a result of a systemic disturbance, such as recurrent blood loss, the underlying cause or causes should be corrected, if possible.

Folic Acid: Folic acid in doses above 1.0 mg daily may obscure pernicious anemia in that hematologic remission can occur while neurological manifestations remain progressive. Pernicious anemia should be excluded before using this product since folic acid may mask the symptoms of pernicious anemia.

Pediatric Use: Safety and effectiveness in pediatric patients have not been established.

Geriatric Use: Clinical studies on this product have not been performed in subjects aged 65 and over to determine whether elderly subjects respond differently from younger subjects. In general, dose selection for an elderly patient should be cautious.

ADVERSE REACTIONS: Adverse reactions with iron therapy may include constipation, diarrhea, nausea, vomiting, dark stools and abdominal pain. Adverse reactions with iron therapy are usually transient. Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

OVERDOSAGE: The clinical course of acute iron overdose can be variable. Initial symptoms may include abdominal pain, nausea, vomiting, diarrhea, tarry stools, melena, hematemesis, hypotension, tachycardia, metabolic acidosis, hyperglycemia, dehydration, drowsiness, pallor, cyanosis, lassitude, seizures, shock and coma.

DOSAGE AND ADMINISTRATION:

Usual adult dose is 1 tablet daily, or as directed by a physician.

DESCRIPTION: CORVITE[®]150 are red, oval bisected tablets embossed "VP052."

HOW SUPPLIED: CORVITE[®]150 tablets are supplied in bottles of 30 units.

Product Code: 68025-052-30

PATENT PENDING

STORAGE: Store at controlled room temperature 15°-30°C (59°-86°F) [see USP]. Protect from light and moisture and avoid excessive heat. Dispense in a tight, light resistant container as defined by the USP using a child-resistant closure.

KEEP OUT OF THE REACH OF CHILDREN.

Rx ONLY

Call your doctor about side effects. You may report side effects by calling: 1-877-95-VERTI (1-877-958-3784).

** Quatrefolic[®] is a registered trademark of Gnosis, SpA.

U.S. Patent No. 7,947,662.

Manufactured for:



Vertical Pharmaceuticals, Inc.

Sayreville, NJ 08872

www.verticalpharma.com

Rev. 08/13



3 68025-052-30 6

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. **KEEP THIS PRODUCT OUT OF REACH OF CHILDREN.** In case of accidental overdose, call a doctor or poison control center immediately.

DOSAGE & ADMINISTRATION
Usual adult dose is 1 tablet daily,
or as directed by a physician.

STORAGE
Store at controlled room temperature 15°-30°C (59°-86°F) [see USP]. Protect from light and moisture and avoid excessive heat. Dispense in a tight, light resistant container as defined by the USP using a child-resistant closure.

Left Margin
Product
Information