

73317-1150-9



PrenatVite^{Plus}

**RX Prenatal Vitamin Tablet with
Prescription Folic Acid**

PRENATAL / POSTNATAL

**R_x Only
Dietary Supplement
90 Tablets**

Each prenatal tablet contains the following ingredients
for nutritional supplement :

Vitamin A (As Acetate and Beta Carotene)	1200	MCG
Vitamin C (Ascorbic Acid)	120	MG
Vitamin D3 (As Cholecalciferol)	10	MCG
Vitamin E (As DI-Alpha Tocopheryl Acetate)	9.2	MG
Vitamin B1 (As Thiamine Mononitrate)	1.84	MG
Vitamin B2 (Riboflavin)	3	MG
Vitamin B3 (As Niacinamide)	20	MG
Vitamin B6 (As Pyridoxine Hcl)	10	MG
Folic Acid	1000	MCG
Vitamin B12 (As Cyanocobalamin) On DCP	12	MCG
Calcium (Calcium Carbonate)	200	MG
Iron (As Ferrous Fumarate)	27	MG
Zinc (As Zinc Oxide)	25	MG
Copper (As Cupric Oxide)	2	MG
Inositol	5	MG
Citrus Bioflavonoid Complex 50%	2	MG
Boron (As Boron Citrate)	0.2	MG

OTHER INGREDIENTS

MICRO CRYSTALLINE CELLULOSE, STEARIC ACID, MAGNESIUM STEARATE,
CROSCARMELOLOSE SODIUM, SILICON DIOXIDE, HPMC E15,
HPMC E5/E6, PEG - 8000, TALC, FD&C RED # 40 (LAKE),
FD&C BLUE # 2 (LAKE), FD&C YELLOW # 5 (LAKE).

Description: Prenatvite PLUS Tablet is an orally administered prenatal/postnatal prescription folic acid-containing dietary supplement and should be administered under the supervision of a licensed medical practitioner.

DOSAGE AND ADMINISTRATION: Before, during and/or after pregnancy, one tablet daily, with a meal, or as directed by a licensed healthcare practitioner regardless of lactation status.

See package insert for full prescribing information.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

CAUTION: High levels of folic acid may, especially in older adults, hide signs of Vitamin B-12 deficiency (such as pernicious anemia), a condition that can cause nerve damage.

Store at 15-30°C (59-86°F). [See USP Controlled Room Temperature.] Protect from light and moisture. Dispense in a tight, light-resistant container. Notice: Contact with moisture may produce surface discoloration or erosion.

Call your licensed medical practitioner about side effect.

You may report side effects by calling AUM 1-631-524-5758.

Manufactured & Distributed By :

AUM Pharmaceuticals
Hauppauge, NY 11788.
Made in USA



N
3

73317 11509 6