SIMPLY FORMULATED TO DELIVER MORE.

The Benefits Line™ is designed for power. Each formula offers clinically relevant levels of evidence-based nutrients to provide more with less.

Keep out of reach of children. Store in a cool, dry place.







LycoBeads® is a registered trademark of LycoRed Ltd Meriva® & Phytosome™ are registered trademarks of Indena SpA. BroccoPhane® is a registered trademark of Cyvex Nutrition, Inc.

*This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

Distributed by DaVinci® Laboratories of Vermont a division of FoodScience® Corp.

929 Harvest Lane Williston, VT 05495 USA www.davincilabs.com To obtain product information or report a serious adverse event, call 1-800-325-1776.



DaVinci[®] BENEFITS

LINE

PHYTO BENEFITS™

A Phytonutrient Supplement for Antioxidant Protection*

VEGETARIAN / GLUTEN FREE / SOY FREE

60 CAPSULES



Suggested Use: As a dietary supplement, adults take one capsule daily, or as recommended by your healthcare practitioner.

Supplement Facts

Serving Size 1 Capsule Servings Per Container 60

Amount Per Serving

% Daily Value

BroccoPhane® Broccoli Sprout Powder 100 mg (standardized to 4,000 ppm Sulforaphane)

Green Tea (Camellia sinensis L.) Leaf Extract 100 mg (Standardized to 70% EGCG)

Meriva® Curcumin Phytosome™ (Curcuma longa root extract/ Phosphatidylcholine complex) 100 mg

Pomegranate (Punica granatum) Fruit Extract 100 mg

(Standardized to 40% Punicosides) Quercetin Dihydrate 100 mg

LycoBeads® natural tomato extract 50 mg

containing [5%] 2.5 mg Natural Lycopene Polygonum cuspidatum Root Extract 50 mg

(Standardized to 50% trans-resveratrol)

*Daily Value not established

Other ingredients: hypromellose (capsule), microcrystalline cellulose, vegetarian leucine.

Color of this product may vary due to color variations of the ingredients

Warning: If pregnant or nursing, consult your healthcare practitioner before taking this product.

Caution: Discontinue use 2 weeks prior to surgery.

0418R