WHEN IT'S TIME TO GO TO BATTLE, YOU GO HARD WITH TOTAL WAR® EXPERIENCE A WORKOUT WITH INTENSITY YOU DIDN'T THINK WAS POSSIBLE. NEW FOUND STRENGTH AND ENERGY ARE JUST MINUTES AWAY. LOCK-IN AND GET READY TO BRING THE NOISE!"*



TOTAL WAR SUGGESTED USE:

WHEN IT COMES TIME TO BRING THE NOISE" FOR A WORKOUT. ASSESS YOUR TOLERANCE BY TAKING (1/2) SCOOP PRIOR TO TRAINING. ONCE YOUR TOLERANCE IS ASSESSED MIX (1) SCOOP WITH 4-6 OUNCES OF WATER 30 MINUTES PRIOR TO TRAINING. DO NOT EXCEED (1) SCOOP IN A 24-HOUR PERIOD.

These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.



Supplement Facts Serving Size: 1 Scoop (14.7g) Servings Per Container: 30

Amount Per Serving	
Citrulline Malate (2:1 Ratio of L-Citrulline and Malic Acid)	6 ç
Beta-Alanine	3.2g
Agmatine Sulfate	Ιg
Taurine	10
Caffeine Anhydrous	250mg
Juniper (Juniperus communis) (berry) Extract	150mg
Di-Caffeine Malate	100mg
Green Tea (leaf) Extract (providing 90% polyphenols)	45mg
Naringin (Citrus grandis) (fruit) (providing 98% Naringin)	25mg
BioPerine® (Black Pepper Fruit Extract)	10mg
Theobroma Cocoa (seed) Extract (providing 98% Theobromine)	5mg

*Daily Value not established

Other Ingredients: Natural and artificial flavors, Disodium phosphate, Silica, Calcium silicate. Sucralose, Citric acid. Maltodextrin, Acesulfame-potassium

DISTRIBUTED BY REDCON1, LLC 701 Park of Commerce Blvd. Boca Raton, FL 33487

1(855)733-2661

WARNING: This product is intended to be consumed by healthy adults 18 years of age or alder. This product contains 350mg of Coffeino per serving. Do not use if you are program, nursing, sensitive to coffeine or other stimulants, kidney disease, seizure diserder, thyroid disease, psychiatric disease, diabetes, prostate enlargement or if you are taking a MAD Inhibitor. Do not use if you are prone to dehydration or exposed to excessive heat. As with any dietary supplement, consult a bealthcare professional before using this product. Discontinue 2 weeks prior to surgery, Immediately discontinue use and consult a healthcare professional if you experience any objects reactions.

