## THE OBJECTIVE:

WHEN IT'S TIME TO GO TO BATTLE, YOU GO HARD WITH TOTAL WAR® EXPERIENCE A WORKOUT WITH INTENSITY YOU DIDN'T THINK WAS POSSIBLE. NEW FOUND STRENGTH AND ENERGY ARE JUST MINUTES AWAY. LOCK-IN AND GET READY TO BRING THE NOISE!"\*



## TOTAL WAR SUGGESTED USE:

WHEN IT COMES TIME TO BRING THE NOISE" FOR A WORKOUT, ASSESS YOUR TOLERANCE BY TAKING (1/2) SCOOP PRIOR TO TRAINING. ONCE YOUR TOLERANCE IS ASSESSED MIX (1) SCOOP WITH 4-6 OUNCES OF WATER 30 MINUTES PRIOR TO TRAINING. DO NOT EXCEED (1) SCOOP IN A 24-HOUR PERIOD.

These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.









**PREWORKOUT** 

Net Wt. 15.54 oz (441a)

## Serving Size: 1 Scoop (14.7g) Servings Per Container: 30

Amount Per Serving Citrulline Malate (2:1 Ratio of L-Citrulline and Malic Acid) Beta-Alanine Aamatine Sulfate Caffeine Anhydrous 250mg Juniper (Juniperus communis) (berry) Extract 150mg 100mg 45mg\* Green Tea (leaf) Extract (providing 90% polyphenols) Naringin (Citrus grandis) (fruit) (providing 98% Naringin) 25mg\* BioPerine® (Black Pepper Fruit Extract) 10mg<sup>3</sup>

\*Daily Value not established

Other Ingredients: Natural and artificial flavors, Disodium phosphate, Silica, Calcium silicate, Sucralose, Citric acid, Maltodextrin, Acesulfame-potassium.

Theobroma Cocoa (seed) Extract (providing 98% Theobromine)

DISTRIBUTED BY REDCON1. LLC 701 Park of Commerce Blvd. Boca Raton, FL 33487

1(855)733-2661

WARNING: This product is intended to be consumed by healthy adults 18 years of age or after. This product contains
350mg of Cofficine per serving. Do not use if you are program, rusting, tensitine to cofficine or other stimulants,
toking any prescription or over-the-counter medications or supplements, or if you have no suspect you may have a
medical condition, including but not limited to, high or low blood pressure, credit or entrytheting, streke, heart, lives or kidaev disease, seizure diserder, thyroid disease, psychiatric disease, diabetes, prostate enlargement or if you are supplement, consult a healthcare professional before using this product. Biscontinue 2 weeks prior to surgery Immediately discontinue use and consult a healthcare professional if you experience any otherse reactions.

