

DIETARY SUPPLEMENT

NIALIPIN is well tolerated by sensitive individuals. Hypersensitive individuals are advised to start with one tablet prior to increasing dosage. Each sustained-release tablet is designed to release niacin over a period of 6 to 8 hours. Do not cut tablets in half.

DIRECTIONS

Take one or two tablets daily, or as directed by a physician.

Do not take niacin unless recommended by and taken under the supervision of a physician if you have any of the following conditions: gallbladder disease, gout, arterial bleeding, glaucoma, diabetes, impaired liver function, peptic ulcer, or if you are pregnant or lactating.

 Manufactured Exclusively For:
CARDIOVASCULAR RESEARCH, LTD.
1061-B Shary Circle, Concord, CA 94518
1-800-888-4585



NIALIPIN
400 mg
Sustained-Release Niacin
Dietary Supplement
60 TABLETS

Supplement Facts

Serving Size: 1 Tablet
Servings Per Container: 60

	Amount Per Serving	% Daily* Value
Niacin (as nicotinic acid)	400 mg	2000%

* Percent Daily Values are Based on a 2,000 Calorie Diet.
** Daily Value Not Established

Other Ingredients: Microcrystalline cellulose, hydroxypropylmethylcellulose, garbanzo bean flour, stearic acid, silicon dioxide, ethylcellulose, food grade.
Gluten free, Casein free

This product has not been evaluated by the FDA. This product is not intended to diagnose, cure, treat or prevent any disease.

