

Supplement Facts

Serving Size: 1 Caplet
Servings Per Box: 100

	Amount per Serving	% DV for Pregnant and Lactating Women
Vitamin A (as Beta Carotene)	630 mcg RAE (2100 IU)	40%
Vitamin C (Ascorbic Acid)	120 mg	100%
Vitamin D (as Cholecalciferol)	8 mcg (32 IU)	50%
Vitamin E (as d-Alpha Tocopheryl Succinate)	13 mg* (20 IU)	60%
Thiamin (as Thiamine Mononitrate)	2 mg	140%
Riboflavin	3.4 mg	200%
Niacin (as Nicotinamide)	10 mg NE	50%
Vitamin B ₆ (as Pyridoxine HCl)	10 mg	500%
Folate	2083 mcg DFE (250 mcg Folic Acid)	347%
Vitamin B ₁₂ (as Cyanocobalamin)	15 mcg	530%
Pantothenic Acid	5 mg	70%
Iron (as Ferrum Pyrophosphate)	50 mg	85%
Magnesium (as Magnesium Oxide)	15 mg	4%
Zinc (as Zinc Oxide)	10 mg	77%
Copper (as Copper Gluconate)	1 mg	77%
*Equal to mcg alpha-tocopherol		

Other Ingredients: Microcrystalline Cellulose, Di-Calcium Phosphate, Croscarmellose Sodium, Pink Color Coating (Hydroxypropylmethyl Cellulose, Polyvinyl Alcohol, Titanium Dioxide, Polyethylene Glycol, Ink, FD&C Red #10 Lake), Fumed Silica, Stearic Acid, Magnesium Stearate, Polydioxane R20, Lactose.

GLUTEN, SUGAR, AND LACTOSE-FREE

Bile OB™ is a prescription multivitamin/multimineral indicated for use in improving the nutritional status of women prior to conception, throughout pregnancy, and in the postnatal period for both lactating and non-lactating mothers.

CONTRAINDICATIONS

Bile OB™ should not be used by patients with a known hypersensitivity to any of the listed ingredients.

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. KEEP THIS PRODUCT OUT OF REACH OF CHILDREN. In case of accidental overdose, call a doctor or poison control center immediately.

PRECAUTIONS

Folic acid alone is an improper therapy in the treatment of pernicious anemia and other megaloblastic anemias where vitamin B₁₂ is deficient. Folic acid in doses above 0.1 mg daily may obscure pernicious anemia, in that hematologic remission can occur while neurological manifestations remain progressive. Allergic sensitization has been reported following both oral and parenteral administration of folic acid.

The patient's medical conditions and consumption of other drugs, herbs, and/or supplements should be considered.

DRUG INTERACTIONS

Bile OB™ caplets are not recommended for and should not be given to patients receiving levodopa because the action of levodopa is antagonized by pyridoxine. There is a possibility of increased bleeding due to pyridoxine interaction with anticoagulants (e.g., Aspirin, Heparin, Clopidogrel).

ADVERSE REACTIONS

Adverse reactions have been reported with specific vitamins and minerals but generally at levels substantially higher than those contained herein. However, allergic and idiosyncratic reactions are possible at lower levels. Iron, even at the usual recommended levels, has been associated with gastrointestinal intolerance in some patients.

DESCRIPTION

Bile OB™ is a pink capsule shaped caplet imprinted with "VP 010".

DIRECTIONS FOR USE

Take one caplet by mouth daily or as directed by a physician.

HOW SUPPLIED

Bile OB™ is available in a unit dose pack of 100 (10x10) caplets.

Product Code: 13811-027-10

STORAGE: Store at controlled room temperature 15°-30°C (59°-86°F). [See USP]. Protect from light and moisture and avoid excessive heat.

KEEP THIS PRODUCT OUT OF REACH OF CHILDREN.

For use on the order of a healthcare practitioner

Call your doctor about side effects. To report side effects, call Verical Pharmaceuticals at 1-770-309-4300 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

PLR-E0B-00002-1

Rev. 12/2021

Manufactured for:
Trigen Laboratories, LLC
Alpharetta, GA 30005
www.trigenlab.com



13811-027-10



Elite OB™ is a prescription multivitamin/multimineral indicated for use in improving the nutritional status of women prior to conception, throughout pregnancy, and in the postnatal period for both lactating and nonlactating mothers.



100 CAPLETS

13811-027-10

Prescribed
Multivitamin



Elite OB™ is a prescription multivitamin/multimineral indicated for use in improving the nutritional status of women prior to conception, throughout pregnancy, and in the postnatal period for both lactating and nonlactating mothers.

