

100
TABLETS



DIETARY SUPPLEMENT

Niacin
50 mg



TAMPER RESISTANT - DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

DIRECTIONS: Adults only: One tablet daily as a dietary supplement, or as directed by a physician.

Supplement Facts

Serving Size 1 Tablet

Amount per tablet		% Daily Value
Niacin (Nicotinic Acid)	50 mg	250%

Other Ingredients: Lactose, microcrystalline cellulose, sodium bicarbonate, stearic acid and magnesium silicate. **Contains Milk.**

CAUTION: Niacin may cause temporary flushing, itching and tingling, feelings of warmth and headache, particularly when beginning, increasing dose, or changing brand of niacin. These effects seldom require discontinuing niacin use. Skin rash, upset stomach, and low blood pressure when standing are less common symptoms; if they persist, contact a physician.



Free of starch, gluten, preservatives, yeast, corn, egg, soy, nuts, artificial colors-flavors.

To preserve quality and freshness, keep bottle tightly closed.

Store between 15°-30°C (59°-86°F).

KEEP OUT OF REACH OF CHILDREN



0 0536-4070-01 0

R0907
75

Control No.:

Exp. Date:

Questions or comments?

Call 1-800-645-2158, weekdays 9 am-5 pm ET

Mfd for: **RUGBY LABORATORIES, INC.**

DULUTH, GEORGIA 30097