


DRUG INTERACTION PRECAUTION: Since oral iron products interfere with absorption of oral tetracycline antibiotics, these products should not be taken within two hours of each other.

Questions or comments?
Call **1-800-645-2158**,
weekdays 9 am-5 pm ET

www.rugbylaboratories.com



Store between 15°-30°C (59°-86°F). Do not expose to excessive heat or moisture.



TAMPER RESISTANT - DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

DIRECTIONS: Adults only: One tablet daily, preferably with a meal.

Supplement Facts

Serving Size 1 Tablet

Amount per tablet	% Daily Value	
Iron (as ferrous sulfate)	65 mg	361%
Calcium	20 mg	2%

Other ingredients: dicalcium phosphate, microcrystalline cellulose, croscarmellose sodium, magnesium stearate, FD&C red #40 (Al-lake), FD&C yellow #6 (Al-lake), hypromellose, PEG 400, titanium dioxide

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. **Keep this product out of reach of children.** In case of accidental overdose, call a doctor or poison control center immediately.

WARNINGS: Do not exceed recommended dosage. The treatment of any anemic condition should be under the advice and supervision of a doctor. Occasional gastrointestinal discomfort (such as nausea) may be minimized by taking with meals. Iron-containing medication may occasionally cause constipation or diarrhea.



Lot No.:

Exp. Date:

Mfd for: **RUGBY LABORATORIES, INC.**

DULUTH, GEORGIA 30097

▲ Peel Here

R0612