

PRODUCT BENEFITS: Coenzyme Q-10 is an essential component of the mitochondrial electron transport chain, vital for the production of ATP required for virtually every cellular process.* As a lipophilic antioxidant, Coenzyme Q-10 works to scavenge free radicals that may lead to oxidative stress.*

Statin medications can reduce Coenzyme Q-10 levels in the body. Coenzyme Q-10 can replenish what Statin medications may deplete. Note: Coenzyme Q-10 is not intended to serve as a replacement for Statin therapy, nor should you discontinue taking any prescribed medications while supplementing with Coenzyme Q-10.

*These statements have not been evaluated by the FDA. This product is not intended to diagnose, treat, cure or prevent any disease.

Laboratory Tested to Guarantee Quality
For Health Care Professional Use Only ©2013 PhysiLogics, LLC

PROD. NO. 55167

To refill please call 1-800-765-6775

or visit www.physilogics.com

Carefully Manufactured by PhysiLogics, LLC
2100 Smithtown Ave., Ronkonkoma, NY 11779 U.S.A.



B70931

06F

Physi**o**Logics®

Coenzyme Q-10 30 mg

**Supports Cardiovascular
and Antioxidant Health***

**Dietary Supplement
120 Vegetarian Capsules**

Supplement Facts

Serving Size 1 Capsule

Amount Per Serving	%Daily Value
Coenzyme Q-10 (as Ubiquinone)	30 mg **

**Daily Value not established.

Other Ingredients: Cellulose (Plant Origin), Hypromellose.
Contains <2% of: Silica, Vegetable Magnesium Stearate.

FREE OF: artificial color, artificial flavor, artificial sweetener, preservatives, sugar, starch, milk, lactose, soy, gluten, wheat, yeast, fish, sodium.

Directions: For adults, take one (1) capsule two times daily, preferably with meals or follow the advice of your health care professional. As a reminder, discuss the supplements and medications you take with your health care providers.

TAMPER RESISTANT: Do not use if seal under cap is broken or missing.
KEEP OUT OF REACH OF CHILDREN.

Store at room temperature and avoid excessive heat.

WARNING: If you are pregnant, nursing, taking any medications or have any medical condition, consult your doctor before use. Discontinue use and consult your doctor if any adverse reactions occur.