## MäritzMayer Laboratories

SUGGESTED USE: As a dietary supplement, take one (1) capsule daily. Do not exceed 2 capsules per day.

CAUTION: Do not exceed recommended dose. St. John's Wort may contribute to photosensitivity resulting in skin irritation and redness in persons exposed to strong sunlight or tanning booths. Avoid in patients at risk of bleeding, taking anticoagulants, or with clotting disorders, based on case reports of bleeding. Discontinue use 2-3 weeks prior to some surgical and dental procedures due to increased risk of bleeding. Use cautiously in patients with history of seizure, based on reports of seizure due to Ginkgo seed ingestion. Use cautiously in children. Avoid use in couples who are trying to conceive, based on theoretical reduction of fertility. Pregnant or nursing mothers, children under 18, individuals taking MAO Inhibiting Drugs, or with a known medical condition should consult a physician before using this or any dietary supplement. This product is manufactured and packaged in a facility which may also process milk, soy, wheat, egg, peanuts, fish and crustacean shellfish.

These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

Manufactured Exclusively For: MäritzMayer Laboratories 1.800.432.9130 www.maritzmayer.com



Supplement Facts Serving Size 1 Capsule	
Amount Per Serving	
Ginkgo Biloba (as 24% extract) Phosphatidylserine Complex Acetyl-L Carnitine St. John's Wort (as .3% extract) Glutamine (as L-glutamine hydrochloride) DMAE Bitartrate Bacopin (20% bacosides) Vinpocetine	50mg* 125mg* 50mg* 250mg* 150mg* 50mg* 100mg* 2mg*

Other Ingredients: Silica, magnesium stearate, gelatin.

KEEP OUT OF REACH OF CHILDREN. DO NOT USE IF SAFETY SEAL IS DAMAGED OR MISSING.

STORE IN A COOL, DRY PLACE.

Lot#: 32004

Manufacture Date: 04/2009 Best if used by: 04/2011

3319-5328-5328-30

**60 Capsules**